

## New species of the genus *Filipinmulciber* Vives, 2009 (Coleoptera: Cerambycidae) from the Philippines

Arvīds Barševskis

Barševskis A. 2016. New species of the genus *Filipinmulciber* Vives, 2009 (Coleoptera: Cerambycidae) from the Philippines. *Baltic J. Coleopterol.*, 16(1): 63 - 68.

New species *Filipinmulciber vivesi* sp. n. (Coleoptera: Cerambycidae) from the Luzon Island (Philippines) are described and illustrated. An updated check-list of the genus *Filipinmulciber* is proposed. The genus *Filipinmulciber* in the world fauna is now represented by three species.

Key words: Coleoptera, Cerambycidae, Lamiinae, Pteropliini, Acronia, new species, fauna, Philippines

*Arvīds Barševskis. Coleopterological Research Center, Institute of Life Sciences and Technology, Daugavpils University, Vienības Str. 13, Daugavpils, LV-5401, Latvia. E-mail: arvids.barsevskis@du.lv*

### INTRODUCTION

The genus *Filipinmulciber* Vives, 2009 (Coleoptera: Cerambycidae) belongs to the tribe Homonoeini Thomson, 1864 of the subfamily Lamiinae Latreille, 1825.

The genus *Filipinmulciber* was compared by Vives (2009) with closely related genera *Metamulciber* Breuning and *Mulciber* Pascoe. The genus has been described on the basis of one new species, *F. breuningi* Vives, 2009 from Luzon Island. Later, Vives (2015) described second species of the genus, *F. palawanus* Vives, 2015 from Palawan Island.

In this paper third new species of the genus (and second known species from Luzon Island) is described and illustrated, and updated check-list for the genus is proposed.

### MATERIAL AND METHODS

New species deposited in the Beetles Collection of Coleopterological Research Center, Institute of Life Sciences and Technology, Daugavpils University, Coleopterological Research Centre (Ilgas, Daugavpils Distr., Latvia) (DUBC). Both known specimens of new species have been collected in the Philippines by the local collector.

The laboratory research and measurements have been performed using *Nikon AZ100*, *Nikon SMZ745T* and *Zeiss Stereo Lumar V12* digital stereomicroscopes, NIS-Elements 6D software, and *Canon 60D* and *Canon 1 Ds Mark II* cameras. The map of the Philippine archipelago (Fig. 2) have been drawn using the software *ArcGis 10*.

## RESULTS

### *Filipinmulciber vivesi* sp. nov.

(Fig. 1)

**Type material. Holotype:** Male: Philippines: Luzon Island, Ilocos Norte, Pagudpod, 04.2016. (Fig. 1A).

**Paratype:** Male: same data as the holotype. (Fig. 1B).

**General distribution:** Philippines: Luzon Isl. (Fig. 1E).

**Description.** Body elongate, black, lustrous, surface with black pubescence and numerous spots of white pubescence (Fig. 1A). Ventral side of body black, with white pubescence. Length: 9.9 - 10.9 mm, maximal width of elytra: 3.1 - 3.3 mm.

Head large and flat, transverse, slightly trapezoidal. Eyes slightly convex, prominent, with large facetes. Cheeks slightly extended and covered with pale sparse pubescence. Surface of head shiny, with sparse and coarse punctation, interspaces between punctures with very thin micropunctures. Middle portion of head without visible longitudinal line. Head with two longitudinal bands of white pubescence, which begins at anterior margin of pronotum and ends at basal part of antenna. Labrum dark brown, curved, pubescent, with punctation, with sparse dark pubescence and long setae on anterior margin. Clypeus brown, curved, narrow, transverse, shiny, with delicate pubescence. Mandible shiny, massive, relatively

wide and sharp. Antennae slender, black or dark, with relatively short pale 1<sup>st</sup> - 3<sup>rd</sup> antennomeres; relatively long 1<sup>st</sup> antennomere thickened, with very thin punctures between pubescence. Underside of head smooth.

Pronotum subsquare, smooth and shiny, with sparse and coarse dorsal punctures, with two large rounded or slightly transverse basal spots, between which with one smaller spot of white pubescence; anterior margin finely bordered; basal angles invisible; lateral parts of pronotum with doubled tooth. Scutellum very small, apically rounded.

Elytra elongate, black, glossy, with well developed shoulders, partly covered with dark pubescence, each elytron with eight or more white spots; base of elytra between shoulders and scutellum smooth and shiny; apical portion of elytra along suture with narrow flat keel-shaped elevation; apex of elytra rounded.

Legs black or red-brown, relatively short, slightly shiny, covered with pale pubescence; femora weakly dilated in medial portion; tibia broadened apically, with dense dark and pale pubescence; tarsomeres black or black-brown, covered with pale pubescence dorsally.

*Aedeagus* relatively long, slightly curved, with sharp lamella (Fig. 1C). Parameras (Fig. 1D).

Female unknown.

**Differential diagnosis.** The new species is closely related with *F. breuningi* (Fig. 2), but differs in the shape of the surface drawings: spots are on the head, at the base of pronotum and particularly on the elytra, where spots are much visible or on the frontal portions does not forming a broad transverse band. Besides that, the surface of *F. breuningi* without white spots or forming transverse band or wide spot on basal part of elytra, and with short transverse band and some small spots in apical part. Pronotum of *F. breuningi* without white spots, while pronotum of the new species has three white spots.




Fig. 1. *Filipinmulciber vivesi* sp.n.: A - Holotype, B - Paratype, C - Aedeagus of holotype, D - Parameras, E - map of distribution.




Fig. 2. *Filipinmulciber breuningi* Vives, 2009: A - D - four colour forms of imago, E - aedeagus, F - paramera

*Aedeagus* of the new species more curved, thinner and slender than *F. breuningi*.

**Etymology.** The new species is named after the prominent Spanish entomologist Eduard Vives, specialist on the fauna of Cerambycidae of the Philippines and East Asian Region.

#### Check-list of the genus *Filipinmulciber*

*Filipinmulciber* Vives, 2009

1. *Filipinmulciber breuningi* Vives, 2009 - Philippines: Luzon Island
2. *Filipinmulciber palawanus* Vives, 2015 - Philippines: Palawan Island
3. *Filipinmulciber vivesi* sp.n. Barševskis, 2016 - Philippines: Luzon Island



Fig. 3. *Filipinmulciber palawanus* Vives, 2015 (Vives 2015)

#### Determination key of the genus *Filipinmulciber*

- 1(2) Body covered by golden pubescence. Each elytron with two longitudinal ribs. Head triangular. ....  
..... ***F. plawanus* Vives**
- 2 (1) Body covered by dark pubescence, often with spots of white pubescence. Head quadrangular.
- 3(4) Pronotum at the base with three white spots..... ***F. vivesi* Barševskis, sp. n.**
- 4(3) Pronotum at the base without three white spots..... ***F. breuningi* Vives**

#### ACKNOWLEDGEMENTS

I wish to express my gratitude to my colleagues Dr. Alexey Shavrin and Dr. Alexander Anichtchenko (Daugavpils, Latvia) for valuable comments and suggestions on the manuscript. I thank my colleague Anita Rukmane for help in preparation of all photographs of the beetles, Māris Nitcis for help in preparation of the distribution map, Marina Janovska, Kristīna Aksjuta and Anita Rukmane for the laboratory assistance and mounting of specimens.

#### REFERENCES

Vives E. 2009. New or interesting Cerambycidae from the Philippines (Coleoptera,

Cerambycidae) (Part III). Les Cahiers  
Magellanes, 105: 1-20.

*Received: 15.06.2016.*

*Accepted: 30.01.2016*

Vives E. 2015. New or interesting Cerambycidae  
from the Philippines (Coleoptera,  
Cerambycidae, Lamiinae) (Part XII). Boletín de  
la S. E. A., 56 (30.6.2015): 49-60