

Biochemical Activity of Superoxide Dismutase Enzyme in Liver of *Labeo rohita* in River Gomti

Devbrat Mishra¹, Prashant Singh²

How to cite this article:

Devbrat Mishra, Prashant Singh. Biochemical Activity of Superoxide Dismutase Enzyme in Liver of *Labeo rohita* in River Gomti 2020; 7(1):15-19.

Abstract

Most of aquatic animals are dependent on molecular oxygen. Yet all living forms are prone to oxygen toxicity. This oxygen toxicity has been attributed to reactive oxygen metabolites including the Superoxide radicals(O_2^-), Hydrogen peroxide (H_2O_2) and hydroxyl radicals (OH^-). These highly reactive substances can directly or indirectly cause substantial damage to living cells. The enzyme superoxide dismutase is essential for the survival of all respiring and metabolically active cells. Only respiring and metabolically active cells can generate reactive oxygen metabolites and they are controlled at physiological concentration by a repertoire of cellular antioxidant defenses such as Superoxide Dismutase (SOD), Catalase (CAT), Glutathione Peroxidase (GPx), Glucose-6-Phosphate dehydrogenase (G-6-PDH) and Glutathione-S-transferase (GST).

Keywords: CAT; GPx; G-6-PDH; GST; OH; ROMs.

Introduction

The concept of oxygen as a requirement for aerobic life is mentioned in ancient Indian civilization. For example, the Sanskrit word "PRAN VAYU" meaning the "the gas of life" appears frequently in the "VEDAS". For centuries, considered critical to life itself, oxygen remains and will always remain, one of the essential elements. The Biochemical Scientists have envisaged much interest on the role of oxygen derived free radicals in various diseases. "Reactive Oxygen Species (ROS) or Free Radicals" have been implicated in over hundred diseases from arthritis and haemorrhagic shock to AIDS (Southern 1988, Halliwell and Gutteridge 1989). The ROS is also related with plant defence system (Deepmala, 2019). The wide range of diseases implied increased formation of free radicals leading to cell and tissue injury in most, if not all human disease (Halliwell and Gutteridge; 1984 and 1989). The metabolism of oxygen generates reactive intermediates i.e. free radicals (Otto and Moon, 1995). A radical might donate its

Author's Affiliation: ¹Assistant Professor, ²Research Scholar, Department of Zoology, Tilak Dhari Post Graduate College, Jaunpur, Uttar Pradesh 222002, India.

Corresponding Author: Prashant Singh, Research Scholar, Department of Zoology, Tilak Dhari Post Graduate College, Jaunpur, Uttar Pradesh 222002, India.

E-mail: devbrat1003@gmail.com

unpaired electrons to another molecule. It might take an electron from another molecule in order to pair or it might simply join to that molecule.

Superoxide is the best known free radical of all oxygen derived species (Fridovich, 1978) because it is the best intermediate in the sequential univalent reduction of oxygen that leads to formation of H_2O . (Florence, 1990; Harris, 1992; Winkle et al., 2007). The hazardous effects of reactive oxygen species are quite well-known, however their detoxification, which is one of the prerequisite of aerobic life, cannot be overlooked for this reason. The nature has equipped the biosystem with multiple defenses. (Otto and Moon, 1996; Winkle et al., 2008; Mishra et al., 2008). The antioxidant systems responsible for cellular protection against oxidative stress are

diversified as the free radicals themselves (Heffner and Repine, 1989). The Superoxide Dismutase (SOD) catalyzes the dismutation between two moles of superoxide anion to yield one mole of oxidized product (oxygen) and one mole of reduced product (Klug et al., 1972; Babich et al., 1993). This is analogous to the dismutation of hydrogen peroxide to oxygen and water catalyzed by catalase (Mishra et al., 2008; Singh et al., 2009; Mishra et al., 2018; Dar et al., 2019; Abhijith et al., 2012). In rat, mice and fish, the Mn-SOD is localized in mitochondria whereas the Cu-Zn-SOD in cytoplasm. Various studies have been carried out to study the limnological condition and heavy metal pollution in the fresh water bodies and its impact on fish physiology (Prakash et al., 2015a, 2015b, 2015c; Srivastava and Prakash, 2018; Prakash and Verma, 2019a, 2019b, 2019c, 2020a, 2020b, 2020c; Prakash, 2020a, 2020b; Verma and Prakash, 2018, 2019a, 2019b, 2020a) but no study on activity of superoxide dismutase was done.

The experimental fish *Labeo rohita* is most common fresh water non air breathing edible fish. It has antioxidant defense system which utilizes the enzymatic and nonenzymatic mechanisms. It can be expected that fish antioxidant defense mechanisms depend on oxygen consumptions. This antioxidant defense mechanism of fish will be detected by the Cu-Zn-SOD and Mn-SOD activities in liver (metabolic tissue) of *Labeo rohita*.

Materials and Methods

The fish *Labeo rohita* was collected from different sites of Gomti river at district Jaunpur, U.P. India and stocked in earthen container and acclimatized in laboratory conditions. The physico-chemical parameters of water such as temperature, pH, alkalinity and DO were analyzed by following standard methods.

After acclimatization fishes were sacrificed by decapitation and liver tissues were taken out then homogenized and centrifuged. The clear supernatant was taken for biochemical studies. The protein and superoxide dismutase were estimated by the method of Lowry et al. (1954) and Fridovich and Fridovich, (1969), respectively. The superoxide anion were generated in a system comprised of NADH and PMS. The superoxide anion reduce the nitro blue tetrazolium (NBT) forming a blue formazan measured at 560nm optical density. For the assay the tissue homogenate were diluted 1:4 for liver tissues. To find out the amounts of Cu-Zn-SOD and Mn-SOD in tissues, 2mm KCN solution

was added to the mixture to inhibit Cu, Zn SOD, Manganese SOD remain unaffected. (Fridovich, 1974; Nandi and Chatterjee 1988; Crapo et al., 1978; Mishra et al. 2008). The unit of enzyme activity was defined as the amount of enzyme required to inhibit the optical density at 560nm of NBT reduction by 50% in one minute under assay condition. The data in this paper have been presented with mean \pm mean standard error and the statistical significance of difference between control and experimental group was calculated by ANOVA.

Results and Discussion

Table 1: Physico-chemical parameters of water samples collected from different sites of Gomti river.

Sites	Temp. °C	pH	Alkalinity mg/lit	DO mg/lit
Kalichabad	24.0 \pm 1.35	7.3	194 \pm 16.3	11.9 \pm 1.83
Katghara	24.2 \pm 1.89	7.4	190 \pm 17.8	11.3 \pm 1.07
Gularghat	24.2 \pm 1.18	7.5	199 \pm 13.8	7.90 \pm 0.99
Hanuman ghat	23.8 \pm 1.45	7.1	170 \pm 16.8	11.2 \pm 0.93
Achalaghat	23.8 \pm 0.21	7.0	169 \pm 13.6	11.5 \pm 0.54
Surajghat	24.1 \pm 1.20	7.2	171 \pm 13.9	11.9 \pm 1.96

Table 2: Total Cu-Zn SOD and Mn-SOD activities (unit mg-1 protein) in liver of *Labeo rohita* collected from different experimental sites. (Activity expressed as mean \pm SD of 5 observations).

Experimental sites of River Gomti in Jaunpur	Total SOD	Cu-Zn SOD	Mn-SOD
Kalichabad	9.0 \pm 0.381	6.6 \pm 0.158	2.3 \pm 0.223
Katghara	8.4 \pm 0.165	6.1 \pm 0.007	2.1 \pm 0.158
Gularghat	7.2 \pm 0.370	5.6 \pm 0.212	1.4 \pm 0.158
Hanuman ghat	7.1 \pm 0.444	5.8 \pm 0.244	1.3 \pm 0.200
Achalaghat	8.6 \pm 0.527	6.2 \pm 0.254	2.2 \pm 0.273
Surajghat	9.1 \pm 0.316	6.8 \pm 0.158	2.3 \pm 0.158

Table 2a: ANOVA of Cu/ Zn SOD of data Table 2.

	S.S.	df	MS	F	P
Total	6.0	28	-	-	-
Between experimental sites	5.1	5	1.04	27.7	<0.001
Error	0.9	23	0.0373	-	-

Table 2b: ANOVA of Mn-SOD of data Table 2.

	S.S.	df	MS	F	P
Total	5.70	28	-	-	-
Between experimental sites	5.6	5	1.14	21.43	<0.001
Error	1.4	23	0.0539	-	-

The physico-chemical parameters of the water




Fig. 1: Physico-chemical parameters of different sites of Gomtiriver at Jaunpur.




Fig. 2: Cu-Zn SOD & Mn-SOD activities (unit mg-1 protein) in liver of *Labeo rohita* collected from different sites.




Fig 3: ANOVA of Cu/Zn SOD of data.




Fig 4: ANOVA of Mn-SOD of data.

samples collected from six different sites of Gomti river at Jaunpur. In these experimental sites temperature was almost identical (23.8-24.020C). There was a nominal difference in pH and alkalinity in all the sites but dissolve oxygen show variation in different sites (Table 1).

The liver plays very important role in metabolism. The Superoxide dismutase, Cu-Zn-SOD Mn-SOD activities in liver of *Labeo rohita* are presented in table and Graph 2, Table 2a and Table 2b, respectively. The highest SOD activities was seen in fish collected from experimental site Kalichabad and lowest SOD activities observed in fishes collected from Hanuman ghat (Table and Fig. 2). The highest Cu-Zn-SOD was found in fishes collected from Surajghat site and lowest in Gularghat site (Table and Fig. 2). Similarly the Mn-SOD activity was observed highest and lowest in fishes collected from Kalichabad and Hanuman ghat respectively (Table and Fig. 2). The ANOVA results indicate that SOD activities of liver in *Labeo rohita* collected from different experimental sites were significantly with each other (Table and Fig. 1- 4).

The result shows that the liver of this more active non air breathing fish, *Labeo rohita* possess an enhanced antioxidant defense system while less active fishes have poor antioxidant defense system. More active fish are metabolically active tissue higher O₂ consumption cause oxidative stress which is inhibited by SOD. SOD forms the primary line of defense against oxidative stress.

Conclusion

The present study reveals the role of dissolved oxygen in the activity of superoxide dismutase enzyme in fresh water non air breathing fish *Labeo rohita*. High DO contents produces oxyanion that cause high oxidative activity of SOD. The present study clearly indicated that the physico-chemical parameters affect the SOD, Cu-Zn-SOD and Mn-SOD activities resulting the severe damage of cellular compounds and also conformational changes in nucleic acid, protein, lipids and carbohydrates. The fishes facing such severe problem of reactive oxygen species also produce the ADS (Antioxidant Defense System) to protect themselves.

References

1. Babich H, Palace MR, Stern A. Oxidative stress in fish cells: in Vivo studies. Arch Environ Contam Toxicol 1993;24(2):173-178.

2. Abhijith BD, Ramesh M, RK. Sublethal toxicological evolution of methyl parathione on same hematological and biochemical parameters in Indian major crop Catla-catla. Comp. Clin Pathol 2012;21:55-56.
3. Crapo JD, Me Cord JM, and Fridovich I. Preparation and assay of Superoxide dismutase: Meth. Enzymol 1978;53:382.
4. Dar SKA, Srivastava PP, Varghese T, et al. (2019): Temporal changes in Superoxide dismutase, Catalase and heat shock protein 70 Gene expression, cortisol and antioxidant enzyme activity of *Labeorohita* finger lings subjected to starvation and refeeding. Pub. Med. PMID: 30633944 DOI:10.1016/J.Gene 2018-12-058.
5. Florence TM. Free radicals, antioxidants and anticancer prevention. Proc. Nutr. Soc. Aust. Annu. Conf 1990;15:88-93.
6. Fridovich I. Superoxide dismutases. Adv. Enzymol 1974;41:35-97.
7. Fridovich I. The biology of oxygen radicals. Sciences 1978;201:875-880.
8. Halliwell B, Gutteridge JMC. Lipid peroxidation oxygen radicals, cell damage and antioxidant therapy. Lancet 1984;1:1396-1398.
9. Halliwell B, Gutteridge JMC (1989). Free radicals in biology and medicine 2nd edition Oxford. Clarendon Press.
10. Harris ED. Regulation of antioxidant enzymes FASEB. J 1992;6:2675-2683.
11. Heffner JE and Repine JE. Pulmonary strategies of antioxidant defense. Am. Rev. Respir. Dis 1989;140:531-554.
12. Klug D, Rabani J, Fridovich I. A direct demonstration of the catalytic action of SOD through use of pulse radiolysis. J. Biol. Chem 1972;247:4839-42.
13. Lowry OH, Rosebrogh NJ, Farr AL. et al. Protein measurement with the Folin Phenol reagent. J. Biol. Chem 1954;193:265:268.
14. Mc Cord JM and Fridovich I. Superoxide dismutase. An enzymatic function for erthrocyte (hemocuprein). J. Biol. Chem 1969;244:6049-55.
15. Mishra Devbrat, Rajesh, Winkle, Singh VB. Distribution of antioxidant enzyme SOD in Gomti and Ganga river in liver and skin of *Cirrhinus mrigala* (Ham). Asian J. Env. Sci. 2008;3(2):128-130.
16. Mishra DB, Dubey JK, Singh Prashant, et al. Observation of biochemical property in brain tissue of fish *Channa punctatus* and exposed deltamethrin. IJSR BS 2018;5(5):66-69.
17. Nandi A, Chatterjee IB. Assay of superoxide dismutase activity in animal tissue. J. Bio Sci 1988;13:305-315.

18. Otto DM, Moon TW 3.3:4.4 tetrachloro biphenyl effects on antioxidant enzymes and glutathione status in different tissues of Rainbow Trout. *Pharmacol. Toxicol* 1995;77(4):281–287.
19. Prakash S. Fish diversity of Semara Taal, a wetland of district Siddharthnagar (U.P.), India. *International Journal of Fisheries and Aquatic Research* 2020a;5(2):07–09.
20. Prakash S. Conservation status of fishes reported from Semara Taal of District Siddharthnagar (U.P.), India. *Internal Journal of Fauna and Biological Studies* 2020b;7(3):21–24.
21. Prakash S, Verma AK, and Prakash S. Limnological Studies of Alwara Lake of Kaushambi (U.P.). *International Journal on Biological Sciences* 2015a;6(2):141–144.
22. Prakash S, Verma AK. Studies on different fish genera in Alwaralake of Kaushambi. *Bioherald: An International Journal of Biodiversity & Environment* 2015b;5(1-2):60–62.
23. Prakash S, Verma AK and Prakash S. Seasonal variation of Zooplankton and Zoobenthos Population in Alwara lake of District Kaushambi (UP) India. *The Journal of Zoology Studies* 2015c;2(5):13–16.
24. Prakash S, Verma AK. Length-Weight Relationships and Condition Factors of Fresh Water Fishes of Baghel Taal of Bahraich (U.P.). *Journal of Experimental Zoology India* 2019a;22(1):343–345.
25. Prakash S, Verma AK. Impact of Arsenic on lipid metabolism of a fresh water cat fish, *Mystus vittatus*. *Journal of Fisheries and Life Sciences* 2019b;4(1):33–35.
26. Prakash S, Verma AK. Acute toxicity and Behavioural responses in Arsenic exposed *Mystus vittatus* (Bloch.). *International Journal on Agricultural Sciences* 2019c;10(1):1–3.
27. Prakash S, Verma AK. Impact of Arsenic on Protein Metabolism of a fresh water cat fish, *Mystus vittatus*. *Uttar Pradesh Journal of Zoology* 2020a;41(5):16–19.
28. Prakash S, Verma AK. Effect of Arsenic on Serum Biochemical parameters of a fresh water cat fish, *Mystus vittatus*. *International Journal of Biological Innovations* 2020b;2(1):11–19. DOI: <https://doi.org/10.46505/IJBI.2020.2102>.
29. Prakash S, Verma AK. Anomalies in biochemical constituents of Kidney in Arsenic induced *Mystus vittatus*. *Bulletin of Pure and Applied Sciences–Zoology* 2020c;39(1):189–193. DOI: 10.5958/2320–3188.2020.00021.2.
30. Singh AK, Dubey JK, Mishra DB et al. Comparative study of Endosulphan effect on antioxidant enzyme (Cat and GPX) of air breathing fresh water teleost *Clarias* (Linn.) and *H. fossilis* (Bluch) 2009. *The Asian Journal of Animal Science* 2009;4(1):22–25.
31. Singh Deepmala. Allelochemical Stress, ROS and Plant Defence System. *International Journal of Biological Innovations* 2019;1(1):33–35. DOI: <https://doi.org/10.46505/IJBI.2019.1107>.
32. Southern PA. Free radical in medicine II: involvement in human disease. *Mayo. Clin. Proc* 1988;63:390–408.
33. Srivastava N, Prakash S. Morphological, Behavioural and haematological alterations in catfish, *Clarias batrachus* (Linn.) after acute zinc toxicity. *International Journal on Biological Science*. 2018;9(1):72–78.
34. Verma AK, Prakash S. Qualitative and quantitative analysis of macrozoobenthos of Beghel Taal, a wetland of U.P. *Indian Journal of Biology* 2018;5(2):127–130.
35. Verma AK, Prakash S. Impact of Arsenic on Carbohydrate Metabolism of a fresh water cat fish, *Mystus vittatus*. *International Journal on Biological Sciences* 2019a;10(1):17–19.
36. Verma AK, Prakash S. Impact of Arsenic on hematology, condition factor, hepatosomatic and gastrosomatic index of a fresh water cat fish, *Mystus vittatus*. *International Journal on Biological Sciences* 2019b;10(2):49–54.
37. Verma AK, Prakash S. Effect of arsenic on enzyme activity of a fresh water cat fish, *Mystus vittatus*. *International Journal of Fisheries and Aquatic Studies*. 2020;8(3):28–31.
38. Winkle, Mishra DB, Singh VB. Impact of pollution in Gomtiriver on antioxidant defense in gills of *H. fossilis*. *Asian. J. Env. Sci* 2007;2(1 & 2):73–76.
39. Winkle, Mishra D.B. and Singh V.B. (2008). Comparative study of antioxidant enzymes in different tissues of *C. batrachus* (Linn.). *J. Adv. Zool.* 29(2):99–102.