

## Review Article

# Phytochemical, Antidiabetic and Therapeutic Properties of *Zygophyllum*

Zohra Mohammadi<sup>1\*</sup><sup>1</sup>Department of Biology, Faculty of Natural Sciences and Life, University of Mustapha Stambouli, Mascara, Algeria

Received: 13.06.2020; Accepted: 15.09.2019

## Abstract

Natural bioactive compounds, isolated from medicinal plants, have an efficient role in the management of various diseases. They have remained sources of new drugs with diversified structural arrangements and interesting biological activities. Among these medicinal plants, species belonging to the genus *Zygophyllum* are efficient sources of bioactive compounds, mainly antihyperglycemic substances. Hence, they are used in traditional medicine against diabetes. It has been indicated that these species could be useful against the skin, stomach and liver diseases. The present review provides phytochemical profiles and in vitro as well as in vivo biological activities of a number of *Zygophyllum* plants with hypoglycemic properties, available through researches from the selected research articles and books published from 1939 to 2017. The sources were collected from various databases.

**Keywords:** Traditional medicine, Antidiabetic plants, Zygophyllaceae, *Zygophyllum Album*, *Zygophyllum gaetulum*, Triterpenoid saponins

---

\*Corresponding Author: Mohammadi Z. Department of Biology, Faculty of Natural Sciences and Life, University of Mustapha Stambouli, Mascara, Algeria. Email: mdi3zhr@gmail.com.

Please cite this article as: Mohammadi Z. Phytochemical, Antidiabetic and Therapeutic Properties of *Zygophyllum*. Herb. Med. J. 2020;in press.

## Introduction

*Zygophyllum* family includes around 285 species in twenty-two genera, growing in arid and semi-arid, tropical and semi-tropical areas. This family comprises small trees, shrubs, and herbs; usually resinous and some are toxic. The biggest taxonomic group is the subfamily of Zygophylloideae, with near 185 species, distributed into 4 genera, i.e. *Fagonia*, *Zygophyllum*, *Augea*, and *Tetraena* (1-4).

*Zygophyllum* is found in several coastal and inland saline sandy soils of the world, along the Mediterranean seashores and oases, with high tolerance to a salty and dry environment. Species of

this genus are a party of succulent herbs that are resistant against drought. They are salt-tolerant, and live in severe, dry atmospheric conditions. The copiousness of species associated with this genus might be due to their high tolerance to environmental stresses. The growth, expansion, and distribution of *Zygophyllum* species are influence by their dependence on the chemical nature of the soil of their habitats (5-10).

*Zygophyllum* plants have mechanisms to adapt like the buildup of soluble osmolytes for diffusion adjustment, increasing the expression of genes that create channels and anti-porters of cytomembrane to eliminate Na<sup>+</sup> and intracellular accumulation of K<sup>+</sup>, and accretion of



**Figure 1.** Fatty Acids Fraction of *Zygophyllum album* (27).

**Table 1:** Fatty Acids Fraction of *Zygophyllum album* (27).

Constituent	Relative percentage
<i>Octanoic acid</i>	0.69
<i>Decanoic acid</i>	33.46
<i>Dodecanoic acid</i>	5.65
<i>Myristic acid</i>	6.47
<i>Palmitic acid</i>	31.18
<i>Heptadecanoic acid</i>	2.43
<i>Stearic acid</i>	4.51
<i>Linoleic acid</i>	7.32
<i>Arachidic acid</i>	8.29

ascorbate peroxidase and superoxide dismutase (SOD) in order to fight reactive oxygen species (ROS). The presence of salt glands in halophytes prevents the buildup of salt within the herb. Additionally, other mechanisms, like proline accumulation, soluble sugars or increase in the activity of antioxidant enzymes are influential in the adaptation of *Zygophyllum* to salt as the rise in stress causes a remarkable increase in proline and raises the relative salt tolerance (11).

The use of plants for health purposes to cure diseases

and relieve physical suffering has started since a long time ago. Plants have been sources of significant pharmaceuticals. A wide range of species are listed in herbal medicine and aromatherapy. Research for remedies in nature continues to increase, and new biologically active molecules are continuously identified. Moreover, their use as drugs has been proven in laboratories by various biological and clinical tests. *Zygophyllum* is a genus which includes a large number of species, many of which have been used by indigenous populations as remedies to treat



**Figure 2.** *Zygophyllum gaetulum* (35).

certain diseases. Since the identification of diabetes, these populations have been found in these plants as antihyperglycemic treatments.

This paper foregrounds the medicinal significance of some species from the genus *Zygophyllum* and elucidates the phytochemical bioactive compounds responsible for the traditional uses of these species and discusses the pharmacological properties proved experimentally by *in vitro* and *in vivo* analyses. To collect the data, we searched publications in various databases and journals using relevant keywords. Information about *Zygophyllum* was obtained from several books and dissertations. After the documentations, interesting files published from 1939 to 2017 were selected and organized in an electronic repertoire. Subsequently, we extracted all the texts which reported botanical, phytochemical,


ethnobotanical, biological and therapeutic properties.

#### **Systematic and Botanical Description of the Genus *Zygophyllum***

The genus *Zygophyllum* could be a principal genus in the Zygophyllaceae. There are 120 species which is indicative of its abundance in the desert, tropical and semi-tropical, saline and calcareous areas, particularly in Western Asia and Africa. This genus generally contains plants or shrubs having flower buds with unpleasant smell (2,12-13).



**Figure 3.** *Zygophyllum coccineum* (47).



**Figure 4.** Structure of some secondary metabolites identified in *zygophyllum coccineum*.



**Figure 5.** *Zygophyllum simplex* (63).

*Zygophyllum* species are glabrous or nearly glabrous annual plants with branches often fleshy and opposite leaves, membranous or herbaceous petiole, paired or solitary axillary flowers, sepals caducous (4 or 5), orangish-red, white, or yellow petals, sometimes with pale or orange claw along the margin, rarely apetalous. Stamens are frequently twice as many as the petals, inserted at the base of the disc. The Disc is fleshy, habitually uniquely lobed, the fruit is a 4-5-angled loculicidal capsule or an indehiscent capsule with 3-4 longitudinal ridges or wings, or separating into 2-4 dorsally winged mericarps, seeds 1-6 in each loculus, with endosperm (13-14).

It has been indicated that several species of the genus *Zygophyllum* could be used to treat a variety of

diseases. They are also used in traditional medicine in certain regions of the world. *Zygophyllum* species are used as anthelmintic agents and for controlling diabetes mellitus. Many in vitro and in vivo studies have confirmed their biological activities (2,15-17).

#### **Biological Properties**

##### ***Zygophyllum album* L.**

**Description:** *Zygophyllum album* (figure 1) is a perennial low shrub (20 to 30 cm in height) with many-branched stems. It has opposite succulent blue-green leaves with petioles. Its flowers are white, and flowering happens in March. The fruit has 5 dilated lobes welded to their base and then widening the peak (5,9,18-20).

*Zygophyllum album* is found in the whole North

African Sahara, Arabia and tropical east Africa. It is abundant in southern Algeria where it is commonly known as “Agaya” (18,20-21).

mite of date palm *Oligonychus afrasiaticus*. Aqueous extracts from this plant caused a mortality rate of 76% in vivo antibacterial activity against *Bacillus*. The



**Figure 6.** *Zygophyllum fabago* (75).

#### Therapeutic and Biological Activities:


*Zygophyllum album* is used as a diuretic, local anesthetic, antihistaminic, antidiabetic, antispasmodic, antihyperlipidemic, antidiarrheal, anti-inflammatory, carminative, antiseptic and stimulant agent for the treatment of skin diseases, diarrhea, and typhoid (2,21-25).

It has been indicated that this plant could be toxic to the sheep causing high mortality. The toxicity/mortality of the crude plant was observed at 7.5-8.4g.kg<sup>-1</sup>b.w. The oral LD50 and the intraperitoneal LD50 of the alcoholic extract for rats were 5.9g.kg<sup>-1</sup>bw and 2.6g.kg<sup>-1</sup>bw, respectively (26-27).

Several scientific research have revealed interesting activities, such as insecticidal activity against the

extract stops bacillosis after 8 to 10 days of treatment for male rats with non-toxic dose. The herbicidal activity against *Echinochloa crus-galli* was observed. The extract (75 g/L) was a potent inhibitor of seed germination (81.59% inhibition), root growth (98.87% inhibition), and shoot growth (84.61% inhibition) after 4 days of treatment (9,24-25).

The aqueous extract of *Zygophyllum album* exhibited in vivo antihyperglycemic, antihyperlipidemic and antioxidant activities. Consumption of the aqueous extract plant (100 and 300 mg/kg bw) for 15 days led to a noticeable decrease in hepatic glucokinase (GK), glycogen in STZ diabetic mice. Moreover, a remarkable increase in hepatic phosphofructokinase (PFK), and glucose-6 phosphate dehydrogenase (G6PDH) was observed in STZ diabetic mice.



**Figure 7.** Chemical Structures of Zygophylosides O (1), P (2) and Zygocarperoside.

Furthermore, ethanolic extracts of *Zygophyllum album* exhibited in vitro inhibitory activity on pancreatic lipase with IC<sub>50</sub> of 91.07 µg/ml and demonstrated a strong antihyperlipidemic effect (28-29).

In addition, various extracts showed a remarkable degree of ferric reducing power activity ( $7.689 \pm 0.562$  to  $30.177 \pm 2.397$  mM), more efficient inhibition of radical DPPH (IC<sub>50</sub> was ranged from  $22.127 \pm 0.837$  to  $84.104 \pm 3.989$  µg/ml), a powerful anti-acetylcholinesterase activity with IC<sub>50</sub> values of 40-58 µg/ml and the highest β-carotene bleaching capacity. These biological activities depend on the bioactive compounds in each extract type (30-31).

**Phytochemicals:** *Zygophyllum album* contained high contents of saponins (18.47 mg/g dry weight), phenolics (12.94 mg/g– 403.46mg/g) and flavonoids (47.16mg/g) (9, 28). Chemical constituents of *Zygophyllum album* are carbohydrates, proteins/amino acids, tannins, lactones, kaempferol, isorhamnetin, Isorhamnetin-3-O-galactopyranoside, Isorhamnetin-3-O-glucopyranoside, isorhamnetin-3-O-rutinoside, isorhamnetin-3-O-robinobioside, quercetin-3-O-glucoside, alkaloids, such as β-carboline (harmine), n-alkanes (C<sub>12</sub>-C<sub>32</sub>), β-amyrin, stigmasterol, β-sitosterol, fatty acids (Table 1), from

which the main fatty acids were decanoic acid 33.46% and palmitic acid 31.18% (26-27,32-33).

***Zygophyllum gaetulum* Emb. and Maire**

**Description:** *Zygophyllum gaetulum* (figure 2) is a plant with branching small sheets constituted of two fleshy leaflets that could be used as a water reserve.

**Therapeutic and Biological Activities:** *Zygophyllum gaetulum* is extensively used in traditional medicine for its antidiarrheal, anti-eczema, anti-inflammatory, antidiabetic, and antispasmodic properties. It is also used against stomach and liver diseases. Several scientific works have highlighted its hypoglycemia activity (2,16,34-36).



**Figure 8.** *Zygophyllum cornutum* (86).



**Figure 9.** *Zygophyllum geslini* (88).

Berzou *et al.* (37-38) indicated that *Zygophyllum gaetulum* aqueous extract attenuates hypercholesterolemia, oxidative stress in plasma and tissues, and increase plasma paraoxonase 1 (PON 1) activity of hypercholesterolemic rats. *Zygophyllum gaetulum* could have a protective effect against oxidative stress in rats fed with a high-cholesterol diet. A noticeable glutathione reductase (GR) activity was observed in the heart (+14%), brain (+47%), and aorta (+23%). Decreased GSH levels were raised by 78, 35, and 65% in the liver, heart, and aorta respectively. Superoxide dismutase activity was 4- and 1.2-fold higher in the liver and aorta respectively. A high level of catalase (CAT) activity was seen in the liver (+14%), heart (+27%), and aorta (+17%), while glutathione peroxidase (GSH-Px) activity was 2.3 and 1.6 fold higher in the liver and aorta respectively. Plasma total cholesterol and low-density lipoprotein cholesterol concentrations were 1.5- and 2.4-fold lower respectively, whereas there were higher levels of high-density lipoprotein cholesterol (+50%). Moreover, extracts from *Zygophyllum gaetulum* showed, *in vivo*, antipyretic activity (in rats) comparable to acetylsalicylic acid, and exhibited anti-inflammatory activity in mice (39-40).

Skim *et al.* (41) observed that the LD50 infusions was higher than 10g/kg body weight and suggested after oral administration of the plant. This plant has no toxicological effect in rats. However, El Youbi *et al.* (36) who investigated the acute toxicity of aqueous extract in mice by intraperitoneal route

indicated that the determination of a lethal dose 50 is equal to 1.2 g/kg of body weight. The results of their study showed the stimulation of proliferative activity on T lymphocytes.

**Phytochemicals:** The phytochemical analysis revealed that *Zygophyllum gaetulum* contains saponins such as bisdesmosidic triterpene saponins, i.e. zygophyllosides I, zygophyllosides L, zygophyllosides M, and quinovic acid glycosides (42).

***Zygophyllum coccineum* L.**

**Description:** *Zygophyllum coccineum* (figure 3) is a small perennial herb characterized by numerous stems and erect, young, green branches with fleshy leaves and moderately whitish flowers of saline and sandy habitats at the vicinity of the sea. Its flowering time is starting from October to November). It is a plant abundantly found in the Eastern (Arabian) Desert and represents the most widespread *Zygophyllum* species in Egypt and Saudi Arabia, where it is found in distinct habitats and wide soil range. It is also found in the Sindh and Baluchistan provinces of Pakistan (10,43-47).

**Therapeutic and Biological Activities:** Traditional medicinal endorses the use of *Zygophyllum coccineum* in the treatment of rheumatism, gout, cough, asthma, hypertension, flatulent colic and diuresis. The juice from fresh leaves and stems has been utilized as an abrasive cleanser. Furthermore, it has been used in the treatment of skin diseases. It has also been used as both an anthelmintic and antidiabetic agent, and an antipyretic and local anesthetic (2,15,46, 48-51).

*In vivo* studies, monitored on extracts of *Zygophyllum*

*coccineum* exhibited remarkable antihypertensive characteristics in normotensive as well as spontaneously hypertensive rats. The impacts of the aqueous extract of *Zygophyllum coccineum* on rat blood pressure (BP) and on the mesenteric vascular bed were examined in rat model animal. The extract dose-dependently decreased BP and heart rate in normotensive and spontaneously hypertensive rats (SHRs). Moreover, it was able to decrease BP in pithed SHRs. The extract did not have any impact on the basal perfusion pressure of the mesenteric vascular bed. Following the increase in the perfusion pressure with noradrenaline or potassium chloride, the extract dose-dependently reduced perfusion pressure. Nonetheless, in preparations in which the perfusion pressure increased with KCl, the depressor reaction to lower doses of the extract was not observed; while higher doses resulted in responses that were reduced in magnitude in comparison with identical responses in preparations in which the perfusion pressure increased with noradrenaline. According to these authors, extracts of *Z. coccineum* could exhibit remarkable antihypertensive activity that may involve some membrane hyperpolarization (2,52).

High antioxidant activity (90.76%) was observed by DPPH radical assay at 1mg/mL of aqueous methanol extract (51). *Zygophyllum coccineum* has also antimicrobial, larvicidal and cytotoxic properties. 32-77% fungal growth inhibition at a concentration of 30 $\mu$ M was effective against *Phomopsis viticola* and exhibited a moderate activity against *Candida albicans* ATCC 10231, while potent antimicrobial and antifungal activities were shown against *Pseudomonas aeruginosa* and *Fusarium moniliforme*. Moreover, *Zygophyllum coccineum* (ethyl acetate extract) had a significant insecticidal and toxic impact against larvae of *Culex pipiens* L. with 100% mortality after 24 h of treatment (53-55). With regard to the plant toxicity, Al-Salihi and Al-Rammahi (47) reported that acute poisoning diarrhea in a herd of camels due to *Zygophyllum coccineum* is related to a high concentration of saponins. The prominent clinical signs were severe diarrhea, polyuria, and photosensitization.

**Phytochemicals:** Phytochemical compounds (figure 4) isolated from *Zygophyllum coccineum* are ursane-

type triterpene saponins flavonoid glycoside, and sterol glycoside. *Zygophyllum coccineum* contains zygophylloside S, kaempferol-3-rutinoside, Quinovic acid (leaves: 0.36%; fruits: 0.31%; stems: 0.47%), and Zygophyllin (leaves: 28%; fruits 0.26%; stems: 0.18%). The phytochemicals identified in leaf extracts, obtained by acetone solvent, are 1-Hexyl-2-nitrocyclohexane; 2-Octadecyl-propane-1,3-diol; Octadecanal; Cyclohexane,1-(1,5-dimethylhexyl)-4-(4-methylpentyl); 2-Octadecyl-propane-1,3-diol and 2(3h)-Furanone,3-(15-hexadecynylidene) (51,53-54,56).

Amin *et al.* (53) isolated nine ursane-type triterpene saponins. These compounds were examined for their antifungal activity against several significant plant pathogens as well as for their insecticidal effect against two important mosquito species. Among the isolated compounds, five of them indicated 32-77 % fungal growth hindrance at a concentration of 30 $\mu$ M against *Phomopsis viticola*. One of the compounds had 90 % and 80 % mosquitocidal activity at 3.1 $\mu$ g/0.5 $\mu$ L against *Aedes aegypti* and *Culex quinquefasciatus*, respectively.

The contents of metabolites in *Zygophyllum coccineum* were variable with the habitat of the plant. Tannins, alkaloids, glucosinolates, proline, and total ascorbate contents are higher in a desert habitat, while the contents of saponins and total glutathione were higher in coastal than in desert plants (57).

#### ***Zygophyllum simplex* L.**

**Description:** *Zygophyllum simplex* (figure 5) is an annual, perennial plant spreading with up to 40 cm long stems. This glabrous plant has simple leaves, and is sessile, cylindrical to obovate, fleshy up to 15 x 4 mm. It has stipules membranous, filiform, flowers axillary, solitary or in pairs, obovate sepals and yellow petals.

**Therapeutic and Biological Activities:** *Zygophyllum simplex* is used for the treatment of horny patches of skin. Infusion of the leaves acts as a skin cleanser. *Zygophyllum simplex* acts as an anthelmintic, analgesic, anti-inflammatory, hypoglycemic, hypolipidemic and laxative agent. The plant is efficient in the treatment of the cold, flu, and cough. Moreover, it has vermifuge properties and is used as eye drop and to cure heart diseases. Furthermore, the plant showed antibiotic activity and a noticeable

reduction in total serum cholesterol, blood glucose and CK level (2,58-61).

**Phytochemicals:** The plant contains free ascorbic acid, sitosterol glucoside, phenolic acids and flavonoids such as kaempferol, kaempferol-3-O-rutinoside, quercetin, quercetin-3,7-di-O-b-glucopyranoside, isorhamnetin, isorhamnetin-3-O- $\beta$ -D-rutinoside, isorhamnetin-3-O-beta-glucopyranoside, Isorhamnetin-3-O-beta-glucopyranoside-7-O-alpha-rhamnopyranoside, 6''-(2-E-butenoyl) isorhamnetin-3-O-glucoside, Gentisic acid, Gentisic acid 5-O-alpha-rhamnopyranoside, myricitrin, luteolin-7-O- $\beta$ -D-glucoside and quinic acid-3- and alpha-L-rhamnoside (33,59,61,63). The two flavonoids, myricitrin and luteolin-7-O- $\beta$ -D-glucoside have indicated the highest level of antioxidant activity. They were shown to be the most efficient agents in decreasing NF $\kappa$ B p65, IL-6, IL-1 $\beta$  and TNF- $\alpha$  levels at the lowest concentration (1 $\mu$ M), in vivo (63).

#### *Zygophyllum fabago* L.

soil has a high level of salinity. The metal uptake into shoots is moderately small. Hence, this herb may be regarded as a significant choice to re-vegetate neutral pH mine tailings (65-66).

**Therapeutic and Biological Activities:** *Zygophyllum fabago* is employed as an antitussive, expectorant, and anti-inflammatory agent and for relieving pains. Aerial parts at the flowering stages are used as antirheumatic, antihelmintic, cathartic, and antiasthmatic substances. The extract of the plant did not show any toxic sign or mortality at the dose of 5000mg/kg (2,67-70).

*Zygophyllum fabago* extracts have allelopathic effects on *Cuscuta monogyna* Vahl (eastern dodder), a parasitic weed of woody species. The extracts showed higher inhibition impacts on seed germination (up to 90.6%) and on growth (66). The antifungal activities of aqueous extracts on two plant phytopathogenic fungal species (*Fusarium oxysporum* f.sp. *melonis* and *Pythium aphanidermatum*) were investigated by Dana *et al.* (71). The plant extracts repressed the growth of *Fusarium oxysporum* (42.9%) and *Pythium*



**Figure 10.** *Zygophyllum hamiense* (4).

**Description:** *Zygophyllum fabago* (figure 6) is frequently known as Syrian bean-caper, oftentimes found within the arid regions of Southern Europe and within the arid zones of the Mediterranean coastal region, like Afghanistan, Pakistan, Iraq, Iran, Saudi Arabia, Turkey, Spain, France, Italy, and North Africa. It is also found in disturbed sites and waste areas (2,64-66).

This succulent, perennial and multi-branched shrub is characterized by its noticeable ability to adapt to severely polluted neutral mine tailings even when the

*aphanidermatum* (85.3%) at a dose of 10%.

The plant has exhibited a really sturdy activity against the human pathogen yeast; *Candida albicans*. The toxicity of fractions, extracted by hexane, ethyl acetate, chloroform, methanol, butanol and water was specified by flow cytometry on  $2.10^6$  CFU/ml of *Candida albicans* at diverse concentrations of extracts for a period of 1, 2, and 24 h at 25°C. The 1h exposure showed that 1.0 g/ml hexane and ethyl acetate extracts of *Zygophyllum fabago* killed 98% and 90% of cells, respectively. Furthermore, plant extracts might also

have antiviral photosensitizers, but the activities have been observed only at elevated concentrations (2,72-74).

**Phytochemicals:** *Zygophyllum fabago* contains ash (5.43%), moisture (8.26%), amino acids (0.95%), carbohydrates (0.90%), organic acids (7.96%), flavonoids (1.10%), saponins (10.31%), alkaloids (0.44%), and macro/micro-elements (75).

The amino acid and fatty acids profiles of *Zygophyllum fabago* were investigated. The most plentiful essential amino acid and non-essential amino acid were leucine (4.75µg/g) and glutamate (23.42µg/g). The amounts of MUFA (monounsaturated fatty acid) and PUFA (polyunsaturated fatty acid) were 78% and 13.1% of the total fatty acid content, respectively. The monounsaturated fatty acid; Oleic acid; was the main

constituent in *Zygophyllum fabago* (77.1% of total fatty acids), followed by linoleic acid (12.8%). (76).

In the seed oil, the most important constituents were unsaturated fatty acids. The total unsaturated acid was 72.58%. The element with the highest content was linoleic acid (51.02%), followed by oleic acid (21.56%). The amount of palmitic acid (8.34%) was higher than stearic (4.88%), arachidic (1.41%), and behenic acid (1.05%) (69).

The aerial part of the plant contains triterpenoid saponins such as Zygofaboside C, a sulfated saponin with chemical name 3β,23,30-trihydroxyurs-20-en-28-al-23-sulfate 3-O-β-D-xylopyranoside (2,69,77).

The bark of *Zygophyllum fabago* L. contains the 27-nor-triterpenoid glycoside, 3-O-β-d-glucopyranosyl-pyrocincholate, 3-O-6-deoxy-β-d-glucopyranosyl-pyrocincholate, quinovic acid, 3-O-6-deoxy-β-d-

**Table 2:** Contents of Simple Phenols and Flavonoids. Identified in *zygophyllum hamiense* (17).

Constituent	Amount (%)
Pyrogallol	1.94
Gallic acid	2.02
Protocatechuic acid	0.97
Catechin	1.81
Chlorogenic acid	2.31
Catechol	2.57
Caffeic acid	1.54
Vanillic acid	1.37
Ferulic acid	0.84
Salicylic acid	1.10
Benzoic acid	1.51
Coumaric acid	1.20
Chrysin	0.34
Quercetin	0.04
Rosmarinic acid	2.33
Rutin	10.71
Quercitrin	0.77
Naringenin	0.94

glucopyranosyl-quinovic acid, 3-O- $\beta$ -d-glucopyranosyl-quinovic acid, 3-O-6-deoxy- $\beta$ -d-glucopyranosyl-cincholic acid, zygophylosides O and P (figure 7). Compounds such as 3-O- $\beta$ -d-glucopyranosyl-pyrocincholate, 3-O-6-deoxy- $\beta$ -d-glucopyranosyl-pyrocincholate, and quinovic acid have shown certain anti-tumor activities by MTT assay, while Zygophyloside O exhibited weaker nitric oxide (NO) inhibitory activity (67,78).

Phytochemical active compounds identified in the plant roots are zygocaperoside (figure 7) and Isorhamnetin -3-O glucoside (79).

#### ***Zygophyllum cornutum* Coss.**

**Description:** *Zygophyllum cornutum* Coss. (figure 8) is a xerophyte herb known for its dilated fruits on top in a free portion of carpels recurved into hooks as long as the welded portion. It is an endemic plant distributed in the Northern Sahara of Algeria.

**Therapeutic and Biological Activities:** It is used in traditional medicine in the treatment of diabetes, hypertension, dermatitis, rheumatism, gout and asthma. This species has shown hypocholesterolemic, hypolipidemic and antioxidant activities (2,80-82).

It was shown via free radical DPPH assay that crude extract of *Zygophyllum cornutum* and its fractions could have significant antioxidant activities. EC50 ( $\mu\text{g/ml}$ ) enregistered are ( $67.059 \pm 4.727$ ); ( $38.478 \pm 2.085$ ); ( $47.767 \pm 1.571$ ); ( $42.159 \pm 3.029$ ); ( $24.935 \pm 1.983$ ) for crude extract and fractions (chloroform, ethyl acetate, butanol and water) respectively (81). Boumaza (83) has also shown a significant scavenger effect and high antioxidant activity. The results registered by this author revealed increased blood glucose, urea, uric acid, creatinine, low density lipoprotein-cholesterol ( $\text{LDL}_C$ ), cholesterol and triglycerides with decreased level of high-density lipoprotein-cholesterol ( $\text{HDL}_C$ ) and total proteins in diabetic rats. Moreover, TBARS (Thiobarbituric acid reactive substance) in the different organs' tissues were markedly increased while GSH (glutathione) and CAT (Catalase) were significantly modified compared to control normal rats. Oral administration of ZCME (*Zygophyllum cornutum* methanolic extract) in a dose of 700mg/kg.bwt improved serum glucose, lipid profile and renal function tests. TBARS were significantly reduced in all organs'

tissues, while CAT and GSH were markedly restored.

**Phytochemicals:** Phytochemical compounds of *Zygophyllum cornutum* are  $\beta$ -sitosterol, polyphenols, flavonoids such as isorhamnetin-3-rutinoside (narcissin), coumarins, alkaloids, tannins and ursane-type saponins: 3-O-( $\beta$ -D-quinovopyranosyl) quinovic acid, 3-O-(2-O-sulfo- $\beta$ -D-glucopyranosyl) quinovic acid, 3-O-( $\beta$ -D-quinovopyranosyl) quinovic acid-28-(O- $\beta$ -D-glucopyranosyl) ester and its sulfated derivative: zygophyloside E, zygophyloside G, zygophyloside H (83-85).

#### ***Zygophyllum geslinii* Coss.**

**Description:** *Zygophyllum geslinii* Coss. (figure 9) is a species endemic to Algerian central Sahara.

**Therapeutic and Biological Activities:** It is used in traditional medicine as a complement in the treatment of diabetes.

The aqueous and methanol extracts from *Zygophyllum geslinii* were able to hinder the activity of  $\alpha$ -amylase in proportion to the concentration of the extracts. Extracts from *Zygophyllum geslinii* are equipped with an important antioxidant potential. Activities of 88.22% and 75.89% are noted for a concentration of 2mg/ml with methanolic and aqueous extracts.

**Phytochemicals:** This medicinal plant contains flavonoids, saponins, tannins and alkaloids, glycosides zygophylosides E, G, and H, and 3-O-( $\beta$ -D-quinovopyranosyl) and quinovic acid 28-(O- $\beta$ -D-glucopyranosyl) ester. Total polyphenols and flavonoids values are 7.5mg EGA/g and 2.41mgCEQ/g for the methanolic extract and 6.3mg EGA/g and 2.32 mgCEQ/g for the aqueous extract (87-89).

#### ***Zygophyllum hamiense* Schweinf.**

**Description:** *Zygophyllum hamiense* (figure 10) is a perennial shrubs, with green, reddish or yellowish green colors. Its height and width reach up to 80cm and 90cm respectively. Its stems are pubescent with unicellular simple trichomes. Leaves of this plant are mostly 1-foliolate, sometimes 2-foliolate in upper branches, terete, globular or cylindrical or clavate, 4-9 mm long, 3-6 mm wide, fleshy, pubescent or glabrous, petiole equal or longer than leaflets up to 9 mm long, stipules triangular, herbaceous,  $1 \times 1.5$  mm, pubescent. Flowers are bisexual, solitary at each node, white, 4-6  $\times$  3-5 mm, and pedicle is 3-5 mm long. This plant is found largely along the Persian Gulf region and grows

on salt accumulated lands.

**Therapeutic and Biological Activities:** The plant is commonly used as firewood but its sprouts are used to feed camel. It has been indicated that the extract of this plant is a potent DPPH free radical scavenger. Moreover, the acute toxicity determination of the LD50 revealed that *Zygophyllum hamiense* extracts was found to be safe up to dose 5 g/kg. The maximum flavonoid content (1.48 %) was extracted with acetone, meanwhile ethyl acetate and ethanol was highly beneficial for solubilization of total phenol (4 and 3.61 mg GAE/g dry plant wt., respectively) (4,17).

**Phytochemicals:** *Zygophyllum hamiense* contains polyphenols and flavonoids. Total of 18 phenolic components were identified in *Zygophyllum hamiense* (Table 2), where rutin is the major phenolic constituent (17).

#### *Zygophyllum eurypterum* Boiss. & Buhse

**Description:** *Zygophyllum eurypterum* is an erect shrub. It is 1m to 1.2m tall. Stem and branches are whitish grey. Leaves are 1.5-3 cm long. Flowers are creamy white. Seeds are brown and 6-7 mm long. It is widely distributed in Iran, Afghanistan, Russia and Pakistan (80-91).

**Therapeutic and Biological Activities:** It has been shown that antimicrobial activities are efficient against *Staphylococcus aureus*, *Escherichia coli*, *Aspergillus niger* and *Candida albicans*. Moreover, antioxidant activities examined in the stem and leaves and evaluated by DPPH radical scavenging activity revealed the respective IC50 values of 80.41% and 424.4% for n-hexane fraction and methanolic extracts. The IC50 values are 15.37% and 84.61% for the stem and leaves respectively. On the other hand, the respective % inhibitions in linoleic acid system for the stem and leaves vary from 26.7% (n-hexane) to 96% (methanol) and from 44.7% (n-hexane) to 87.5% (n-butanol) (91).

**Phytochemicals:** *Zygophyllum eurypterum* contains stigmasterol,  $\beta$ -sitosterol,  $\beta$ -sitosterol glucoside, oleanolic acid, harmine, 7,3',4'-trimethoxyflavone, 5-methylflavanone, atriplicoid A, atriplicoid B and four pterocarpan, atricarpan A atricarpan B, atricarpan C and atricarpan D. It was shown that atricarpan D could inhibit AChE with an

IC50=20.5 $\mu$ M.

Total phenolic contents observed in the leaves of this plant ranged from 5.52 mg/100 g (n-hexane) to 98.25 mg/100 g (methanol). Similarly, the range of total flavonoids contents varied from 1.46 mg/100 g (n-hexane) to 24.41 mg/100 g (methanol) in the stem and 0.70 mg/100 g (n-hexane) to 4.46 mg/100g (methanol) for leaves (92-94).

## Conclusion

*Zygophyllum* species are potential sources of highly interesting active compounds used in the treatment of many diseases. These species could be used as raw materials of primary significance for pharmaceutical industries and drug preparation as well as design. They are particularly used as dietary supplements for the management of diabetes.

## Acknowledgements

I would like to thank all the researchers whose works have contributed in various ways to this review study.

## Conflict of Interest

The authors declare that they have no conflict of interest.

## References

1. Beier BA, Chase MW, Thulin M. Phylogenetic relationships and taxonomy of subfamily Zygophylloideae (Zygophyllaceae) based on molecular and morphological data. *Plant Systematics and Evolution*. 2003;240:11-39.
2. Khan SS, Khan A, Khan K, Farooq U, Ahmed A, Zahoor A, et al. A New Ursane Type Sulfated Saponin from *Zygophyllum fabago* Linn. *Rec Nat Prod*. 2014; 8(4):354-9.
3. Christenhusz MJM, Byng JW. The number of known plants species in the world and its annual increase. *Phytotaxa*. 2016;261(3):201-17.
4. Alzahrani DA, Elbokhari EJ. Systematic studies on the Zygophyllaceae of Saudi Arabia: A new variety and new variety combination in *Tetraena*. *Saudi J Biol Sci*. 2017;24(7):1574-9.
5. Täckholm V. *Students Flora of Egypt*. 2nd ed. Cairo: Univ Press. 1974.
6. Boulos L. *Flora of Egypt*. 2. Geraniaveae-Boraginaceae. Cairo: Al Hadara. 2000.
7. Bellstedt DU, van Zyl L, Marais EM, Bytebier B, de Villiers CA, Makwela AM. et al. Phylogenetic relationships, character evolution and biogeography of Southern African members of genus *Zygophyllum* (Zygophyllaceae) based on three plastid regions. *Mol Phylogenet Evol*. 2008;47(3):932-49.
8. Hammad I, Qari SH. Genetic diversity among *Zygophyllum* (Zygophyllaceae) population based on RAPD analysis. *Genet Mol Res*. 2010;9(4):2412-20.

9. El-Amier YA, Abdullah TJ. Allelopathic Effect of Four Wild Species on Germination and Seedling Growth of *Echinochloa crus-galli* (L.) P. Beauv. *Int J Adv Res.* 2014;2(9):287-94.
10. El-Amier YA, El-Shora HM, Hesham M. Ecological Study on *Zygophyllum coccineum* L. in Coastal and Inland Desert of Egypt. *J agric ecol.* 2016;6(4):1-17.
11. Ahmadi S, Ehsanpour AA. On the Effect of Salinity on Proline, Antioxidant Enzymes, and Protein in *Zygophyllum*. *Australian Journal of International Social Research.* 2015;1(3):76-83.
12. Ghafoor A. *Zygophyllaceae*. In: Nasir E, Ali SI, editors. *Flora of West Pakistan*. Karachi: University of Karachi. 1974;1:35.
13. Wilson AJG. *Flora of Australia*, 26: *Meliaceae, Rutaceae, Zygophyllaceae*. Clayton South: CSIRO Publishing. 2013.
14. Shu W. *Zygophyllum Linnaeus, Sp. Fl China.* 2008;11:45-9.
15. Elgamal MH, Shaker KH, Pöllmann K, Seifert K. Triterpenoid saponins from *Zygophyllum* species. *Phytochemistry.* 1995;40(4):1233-6.
16. Jaouhari JT, Lazrek HB, Jana M. The hypoglycemic activity of *Zygophyllum gaetulum* extracts in alloxan-induced hyperglycemic rats. *J Ethnopharmacol.* 2000;69(1):17-20.
17. Shehab NG, Abu-Gharbieh E, Bayoumi FA. Impact of phenolic composition on hepatoprotective and antioxidant effects of four desert medicinal plants. *BMC Complement and Altern Med.* 2015;15(401):1-12.
18. Hosny AI. Revision of genus *Zygophyllum* L., sections *Bipartia* and *Mediterranean* in Egypt and Arabia. Cairo: Cairo University. 1978.
19. Mosallam HAM. Size Structure of *Zygophyllum album* and *Cornulaca monacantha* Populations in Salhyia Area, East of Egypt. *Int J Agri Biol.* 2005;7(3):345-51.
20. Shahba MA. On the Ecophysiology and Seed Germination of *Zygophyllum album* Native to the Western Mediterranean Coastal Habitats in Egypt. *Aust J Basic & Appl Sci.* 2010;4(8):3643-57.
21. Harek Y, Derfouf H, Rouabhi H, Larabi L. Inhibitive Action of Aqueous Extract of *Zygophyllum Album* L. Leaves on the Corrosion of Carbon Steel in Hydrochloric Acid Solution. *Der Pharm Lett.* 2017;9(4):57-68.
22. Ayad R. Research and structural determination of secondary metabolites of the species: *Zygophyllum cornutum* (*Zygophyllaceae*). *Exact Sciences: University of Constantine.* 2008.
23. El Ghouli J, Ghanem-Boughanmi N, Ben-Attia M. Biochemical study on the protective effect of ethanolic extract of *Zygophyllum album* on streptozotocin-induced oxidative stress and toxicity in mice. *Biomed Prev Nutr.* 2011;1(2):79-83.
24. Lakhdari W, Dehliz A, Acheuk F, Soud A, Hammi H, Mlik R, et al. Acaricidal Activity of Aqueous Extracts against the mite of date palm *Oligonychus afrasiaticus* Meg (Acari: Tetranychidae). *J Med Plants Stud.* 2015;3(6):113-7.
25. Belmimoun A, Meddah B, Meddah Tir Touil A, Tou A, Sonnet P. In Vivo Antimicrobial Activity of Methanolic Extract of *Zygophyllum album* against *Bacillus cereus*. *J Appl Environ Biol Sci.* 2016;6(6):13-9.
26. Attia HA, Samar MM. Antidiarrhoeal activity of some Egyptian medicinal plant extracts. *J Ethnopharmacol.* 2004;92(2-3):303-9.
27. Moustafa AMY, Khodair AI, Hammouda FM, Husseiny HA. Phytochemical and Toxicological Studies of *Zygophyllum album* L. *J Pharmacol Toxicol.* 2007;2(3):220-37.
28. Mnafgui K, Hamden K, Ben Salah H, Kchaou M, Nasri M, Slama S, et al. Inhibitory Activities of *Zygophyllum album*: A Natural Weight-Lowering Plant on Key Enzymes in High-Fat Diet-Fed Rats. *eCAM.* 2012;ID 620384:1-9.
29. El Ghouli J, Ben-Attia M. Vasorelaxant Effects of Aqueous Extract of *Zygophyllum Album* and Antihyperglycemic Activities in Streptozotocin-Induced Diabetic Mice. *J Diabetes Metab.* 2014;5(9):1-7.
30. Belguidoum M, Dendougui H, Kendour Z. In vitro antioxidant properties and phenolic contents of *Zygophyllum album* L. from Algeria. *J Chem Pharm Res.* 2015;7(1):510-14.
31. Kchaou M, Ben Salah H, Mhiri R, Allouche N. Anti-oxidant and anti-acetylcholinesterase activities of *Zygophyllum album*. *Bangladesh J Pharmacol.* 2016;11(1):54-62.
32. Shoaib AME. Comparative study of the common Egyptian *Zygophyllum* species viz., *Zygophyllum coccineum* L., *Zygophyllum album* L. and *Zygophyllum simplex* L. Cairo: University of Cairo. 1957.
33. Hussein SR, Marzouk MM, Ibrahim LF, Kawashty SA, Saleh NAM. Flavonoids of *Zygophyllum album* L.f. and *Zygophyllum simplex* L. (*Zygophyllaceae*). *Biochem Syst Ecol.* 2011;39(4-6):778-80.
34. Bellakhdar J, Claisse R, Fleurentin J, Younos C. Repertory of standard herbal drugs in the Moroccan pharmacopoea. *J Ethnopharmacol.* 1991;35(2):123-43.
35. Idlimam A, Lamharrar A, Abdenouri N, Akkad S. Thermodynamic Properties and Moisture Sorption Isotherms of *Argania spinosa* and *Zygophyllum gaetulum*. *J Agron.* 2008;7(1):1-14.
36. El Youbi AH, Bousta D, Ouahidi I, Aarab L. Primary pharmacological screening of an endemic plant from the Southern Morocco (*Tetraena gaetula* [Emb. & Maire] Beier & Thulin). *C R Biologies.* 2010;333(10):736-43.
37. Berzou S, Taleb-Senouci D, Guenzet A, Krouf D. *Zygophyllum gaetulum* attenuates hypercholesterolemia and protects against oxidative stress in rats fed a high-cholesterol diet. *J Exp Integr Med.* 2014;4(4):255-60.
38. Berzou S, Krouf D, Dida N, Taleb-Senouci D, Guenzet A. Effects of *Zygophyllum gaetulum* Aqueous Extract on Erythrocytes Lipid Peroxidation and Paraoxonase 1 Activity in Hypercholesterolemic Rats. *Int J Pharmacognosy and Phytochem Res.* 2016;8(10):1689-94.
39. Rimbau V, Cerdan C, Vila R, Iglesias J. Antiinflammatory Activity of Some Extracts from Plants used in the Traditional Medicine of North-African Countries (II). *Phytother Res.* 1999;13(2):128-32.
40. Larhsini M, Markouk M, Jaouhari JT., Bekkouche K, Lazrek HB, Jana M. The antipyretic activity of some Moroccan medicinal plants. *Phytother Res.* 2002;16(S1):97-8.
41. Skim F, Lazrek BH, El Amri H, Kaaya A, Jana M. Toxicological studies on *Globularia alypum* and *Zygophyllum gaetulum* in rats. *Phytother Res.* 1998;12(8):592-4.
42. Safir O, Fkih-Tetouani S, De Tommasi N, Aquino R. Saponins from *Zygophyllum gaetulum*. *J Nat Prod.* 1998;61(1):130-4.
43. Soliman GL. From *Zygophyllum coccineum* L., *Zygophyllaceae*. *J Chem Soc.* 1939;1939:1760-1.
44. Batanouny KH, Ezzat NH. *Eco-Physiological Studies on Desert plants. I. Autecology of Egyptian Zygophyllum species.* *Oecologia.* 1971;7(2):170-83.
45. Reed JD. Ecological biochemistry of secondary plant compounds in herbivore nutrition. In: Buchanan-Smith JG, Bailey LD, McCaughey P, editors. *Constraints on Animal Production from Forages and Grasslands.* Canada: XVIII International Grassland Congress. 1997;8-19.
46. Qaisar M, Uddin G, Uddin Ahmad V, Rauf, A. Mass Fragmentation Pattern of New *Zygophyllosides* from *Zygophyllum propinquum* Decne. *Middle-East J Sci Res.* 2011;8(2):526-9.
47. Al-Salihi KA., Al-Rammahi HM. Diarrhea Associated with Ingestion of *Zygophyllum coccineum* (Tartir) in Camels in the Al-Najaf Desert in Iraq. *International Journal of Poisonous Plant Research.* 2014;3(1):1-6.
48. Saber AH, El-Moghazy Shoaib AM. *Zygophyllum coccineum*. V. Chemistry of leaf and stem. *J Pharm Sci UAR.* 1960;1:1-6.
49. Saber AH, El-Moghazy Shoaib AM. Triterpenoid saponins

- from the roots of *Zygophyllum* species. *Pharmaceutical Sci UAR*. 1961;1:135.
50. Rizk AM, El-Ghazaly GA. Medicinal and poisonous plants of Qatar. Qatar: Scientific and Applied Research Center. 1995.
51. Abd El Raheim MD, Aftab A, Younis SN, Radwan M. Evaluation of Antimicrobial and Antioxidant activities of *Matricaria recutita*, *Ricinus communis* and *Zygophyllum coccineum* Extracts. *Bull Env Pharmacol Life Sci*. 2016;5(7):30-3.
52. Gibbons S, Oriowo MA. Antihypertensive effect of an aqueous extract of *Zygophyllum coccineum* L. in rats. *Phytoh Res*. 2001;15(1):452-5.
53. Amin E, El-Hawary S, Fathy MM., Mohammed R, Ali Z, Tabanca N, et al. Triterpenoidal Saponins: bioactive secondary metabolites from *Zygophyllum coccineum*. *Planta Med*. 2011;77(5):488-91.
54. Ara I, Bukhari NA, Solaiman D, Bakir MA. Antimicrobial effect of local medicinal plant extracts in the Kingdom of Saudi Arabia and search for their metabolites by gas chromatography-mass spectrometric (GC-MS) analysis. *J Med Plants Res*. 2012;6(45):5688-94.
55. Elbadry MA, Elaasser MM, Elshiekh HH, Sheriff MM. Evaluation of Antimicrobial, Cytotoxic and Larvicidal Activity of *Zygophyllum Coccineum* North Sinai. *Egypt. Med Aromat Plants*. 2015;4(5):1-6.
56. Batanouny KH. *Wild Medicinal Plants in Egypt*. Switzerland: IUCN, The World Conservation Union. 1999.
57. El-Shora HM, El-Amier YA, Awad MH. Comparative Phytochemical Studies on *Zygophyllum coccineum* L. from Different Habitats, Egypt. *British Journal of Applied Science & Technology*. 2016;15(4):1-9.
58. Al-Saleh FS, Ali HH, Mirza M. Chemical constituents of some medicinal plants growing in Bahrain. *Fitoterapia*. 1993;64(3):251-6.
59. Hameed M, Ashraf M, Al-Quriany F, Nawaz I, Ahmad MSA, Younis A, et al. Medicinal of the Cholistan Desert: a review. *Pak J Bot*. 2011;43(S1):39-50.
60. Shah A, Marwat SK, Gohar F, Khan A, Bhatti KH, Amin M, et al. Ethnobotanical Study of Medicinal Plants of Semi-Tribal Area of Makerwal & Gulla Khel (Lying between Khyber Pakhtunkhwa and Punjab Provinces), Pakistan. *Am J Plant Sci*. 2013;4(1):98-116.
61. Daradka HM. Evaluation of Hematological and Biochemical Activity of Ethanolic Extract of *Zygophyllum simplex* Linn. in Wistar Rats. *Pak J Biol Sci*. 2016;19(4):179-8
62. Hassanean HA, Desoky EK. An acylated isorhamnetin glucoside from *Zygophyllum simplex*. *Phytochemistry*. 1992;31(9):3293-94.
63. Abdallah HM, Esmat A. Antioxidant and anti-inflammatory activities of the major phenolics from *Zygophyllum simplex* L. *J Ethnopharmacol*. 2017;205(9):51-6.
64. Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, et al. *Flora Europaea*. 2. Rosaceae to Umbelliferae. Cambridge: Cambridge University Press. 1968.
65. Conesa HM, Schulin R, Nowack B. A Laboratory Study on Revegetation and Metal Uptake in Native Plant Species from Neutral Mine Tailings. *Water Air Soil Pollut*. 2007;183(1):201-12.
66. Ebrahimi L, Hassannejad S. Allelopathic effects of syrian bean caper (*Zygophyllum fabago* L.) on seed germination and seedling growth of eastern dodder (*Cuscuta monogyna* Vahl). *J Bio & Env Sci*. 2015;7(2):253-60.
67. Feng YL, Wu B, Li HR, Li YQ, Xu LZ, Yang SL, Kitanaka S. Triterpenoidal saponins from the barks of *Zygophyllum fabago* L. *Chem Pharm Bull*. 2008;56(6):858-60.
68. Baytop T. *Therapy with medicinal plants in Turkey (Past and Present)*. Istanbul: Nobel Medicine Publications. 1999.
69. Erdemoglu N, Kusmenoglu S. Fatty Acid Composition of *Zygophyllum fabago* Seeds. *Chem Nat Compd*. 2003;39(6):595-96.
70. Alhaddad H, Fadhil AA, Ismael FS. Estimation of LD50 and Acute Toxicity of *Zygophyllum fabago* in Mice. *Am J of Pharmacol Sci*. 2015;3(4):94-7.
71. Dana ED, De Lomas JG, Sánchez J. Effects of the aqueous extracts of *Zygophyllum fabago* on the growth of *Fusarium oxysporum* f. sp. *melonis* and *Pythium aphanidermatum*. *Weed Biol Manag*. 2010;10(3):170-5.
72. Zaidi MA, Crow SAJ. Biologically active traditional medicinal herbs from Balochistan, Pakistan. *J Ethnopharmacol*. 2005;96(1):331-4.
73. Hudson JB, Lee MK, Sener B, Erdemoglu N. Antiviral Activities in Extracts of Turkish Medicinal Plants. *Pharm Biol*. 2000;38(3):171-5.
74. Zaidi MA, Crow SAJ. Cytotoxicity of Four Medicinal Plants of Pakistan. *Pak J Bot*. 2012;44(SI):395-7.
75. Abdel-Hamid RA. Investigation of chemical constituents and biological activities of some native halophytes. Almaty: Al-Faraby Kazakh National University. 2014.
76. Abdel-Hamid RA, Abilov ZA, Sultanova NA, Saitjanova SB, Gemedzhieva NG. Preliminary phytochemical screening of *Zygophyllum fabago*. *Int J Biol Chem*. 2013;6(2):60-4.
77. Attia AA. Triterpenoidal saponins from the aerial parts of *Zygophyllum coccineum* L. and *Zygophyllum fabago* L. *Pharmazie*. 1999;54(12):931-4.
78. Feng YL, Li HR, Xu LZ, Yang SL. 27-nor-triterpenoid glycosides from the barks of *Zygophyllum fabago* L. *J Asian Nat Prod Res*. 2007;9(6-8):505-10.
79. Yaripour S, Delnavazi MR, Asgharian P, Valiyari S, Tavakoli S, Nazemiyeh H. A Survey on Phytochemical Composition and Biological Activity of *Zygophyllum fabago* from Iran. *Adv Pharm Bull*. 2017;7(1):109-14.
80. Quezel P, Santa S. *New flora of Algeria and southern desert regions*. Paris: CNRS. 1963.
81. Belguidoum M, Dendougui H, Kendour Z, Belfar A, Bensaci C, Hadjadj M. Antioxidant activities, phenolic, flavonoid and tannin contents of endemic *Zygophyllum Cornutum* Coss. from Algerian Sahara. *Der Pharma Chemica*. 2015;7(11):312-7.
82. Aclinou P, Abdessemed K, Massiot G, Olivier LM. *Plantes des Aurès* 3. 1. Structure d'un flavonoïde de *Zygophyllum cornutum*. *Plant Med Phytother*. 1988;22:212-8.
83. Boumaza A. Effect of the metanolic extract of *Zygophyllum cornutum* coss. against the oxidative stress associated with diabetes mellitus and related organs. *Nature and Life Sciences: Constantine university*. 2009.
84. Ayad R, Rahai M, Azouzi S, Louaar S, Dendougui H, Akkal S, et al. Phytochemical Investigation of the Endemic Plant *Zygophyllum cornutum*. *Chem Nat Compd*. 2012;48(2):313-4.
85. Bencharif-Betina S, Miyamoto T, Tanaka C, Kabouche Z, Mitaine-Offer AC, Lacaille-Dubois AL. Ursane-Type Saponins from *Zygophyllum cornutum*. *Nat Prod Commun*. 2013;8(5):573-4.
86. Belguidoum M. A phytochemical approach to differentiate between two species of genus *Zygophyllum*. *Mathematics and Matter Sciences: University of Ouargla*. 2012.
87. Smati D, Hammiche V, Nehari H, Alamir B, Merad R. *Zygophyllum Geslini* Coss.: Chemical Investigation Hypoglycemic Activity. *Acta Hort*. 1993;332:243-348.
88. Boudjelthia KW, Hammadi K, Kouidri M, Djebli N. Evaluation of Antidiabetic Activity of Two Plants *Berberis vulgaris* and *Zygophyllum geslini*. *J Phys Chem Biophys*. 2017;7(1):1-5.
89. Smati D, Mitaine Offer AC, Miyamoto T, Hammiche V, Lacaille-Dubois MA. Ursane-Type Triterpene Saponins from *Zygophyllum geslini*. *Helvetica*. 2007;90(4):712-9.
90. Renz P, Reinhold K. Biosynthesis of 5, 6-dimethylbenzimidazole. *Angew Chem Int Ed*. 1967;6:1083.
91. Shahzad F, Zubair M, Rizwan K, Rassol N, Tareen RB, Riaz

M, et al. Evaluation of biological activities and gc-ms analysis of Zygophyllum eurypterum. Oxid Commun. 2014;38(4A):1978-94.  
92. Zarbad S, Uddin AV, Mohammed KK. Chemical Constituents of Zygophyllum Eurypterum. United States: Lambert Academic Publishing. 2012.  
93. Ahmad VU, Iqbal S, Nawaz SA, Choudhary MI, Farooq U,

Ali ST, et al. Isolation of Four New Pterocarpanes from Zygophyllum eurypterum (Syn. Z. atriplicoides) with Enzyme-Inhibition Properties. Chem Biodivers. 2006;3(9):996-1003.  
94. Iqbal S, Khan A, Ahmad VU, Khan MA, Bader S, Farooq U, et al. Two new triterpenoids from Zygophyllum eurypterum. Nat Prod Commun. 2011;6(2):179-82.

© **Zohra Mohammadi**. Originally published in the Herbal Medicines Journal (<http://www.hmj.lums.ac.ir>), 03.02.2021. This article is an open access article under the terms of Creative Commons Attribution License, (<https://creativecommons.org/licenses/by/4.0/>), the license permits unlimited use, distribution, and reproduction in any medium, provided the original work is properly cited in the Herbal Medicines Journal. The complete bibliographic information, a link to the original publication on <http://www.hmj.lums.ac.ir/>, as well as this copyright and license information must be included