



Contents lists available at UGC-CARE

International Journal of Pharmaceutical Sciences and Drug Research

[ISSN: 0975-248X; CODEN (USA): IJPSPP]

journal home page : <http://ijpsdr.com/index.php/ijpsdr>

Research Article

Evaluation of *in-vitro* Anti-inflammatory Activity of Sinapic Acid

Sathuluri Vineela¹, Chinta Manga Devi², Santh Rani Thakur^{2*}¹Chebrolu Hanumaiah Institute of Pharmaceutical Sciences, Chowdavaram, Guntur- 522019, Andhra Pradesh, India²Division of Pharmacology, Institute of Pharmaceutical Technology, Sri Padmavathi Mahila Visvavidyalayam, Tirupati- 517502, Andhra Pradesh, India

ARTICLE INFO

Article history:

Received: 05 October, 2020

Revised: 10 December, 2020

Accepted: 30 December, 2020

Published: 30 January, 2021

Keywords:

Albumin denaturation, Anti-inflammatory, Heat induced and hypotonicity induced hemolysis, Sinapic acid.

DOI:

10.25004/IJPSDR.2021.130105

ABSTRACT

Sinapic acid (SA) is a potent nutraceutical or natural bioactive compound which is commonly present in the human diet. It was evaluated for its anti-inflammatory activity by *in-vitro* methods like albumin denaturation assay, heat-induced and hypotonicity induced haemolysis, Cyclooxygenase-2 (COX-2), and lipoxygenase (5-LOX) inhibition assay at different concentrations. Aspirin (200 µg/mL) and Indomethacin (100 µg/mL) were used as standard drugs. The results showed that Sinapic acid at a concentration range of 10-500 µg/mL significantly ($p < 0.001$) protected the heat-induced protein denaturation. Heat and hypotonicity induced haemolysis of erythrocyte was significantly ($p < 0.001$) inhibited at the concentration of 500 µg/mL. The (COX-2) and (5-LOX) enzymes were significantly ($p < 0.001$) inhibited by Sinapic acid. The results indicate that Sinapic acid may be a potential anti-inflammatory agent.

INTRODUCTION

Inflammation, a palliate response of the body to various offenses, is a double-edged sword. While it is typically recognized as a process for the remission of diseases, this process's persistence may lead to various diseases associated with chronic inflammation,^[1,2] including arthritis, atherosclerosis, neuropathic pain and even cancer.^[3-5]

Steroids are potent anti-inflammatory agents and have a wide range of adverse effects that increase blood sugar levels, retain fluid, impairs wound healing, increases appetite, and produces heart burn, irritability, and osteoporosis.^[6] Non-steroidal anti-inflammatory agents (NSAIDs) are the most widely used over the counter drugs and have common unwanted effects like gastrointestinal irritability, gastric ulcers, nausea, vomiting, bloating

and nephrotoxicity.^[7] There is a pressing need for the development of novel anti-inflammatory drugs.

Sinapic acid is a phytochemical found in various edible plant spices such as, citrus and berry fruits, vegetables^[8-10] cereals, and oilseed crops.^[11,12] Sinapic acid is reported to be effective against infections,^[13] oxidative stress,^[14] cancer,^[15] diabetes,^[16] neurodegeneration,^[17] and anxiety.^[18] The present study is a preliminary work carried out to assess the anti-inflammatory activity of Sinapic acid.

MATERIALS AND METHODS

Materials

Sinapic acid was purchased from Sigma Aldrich, USA. All the other chemicals used were of analytical grade.

*Corresponding Author: Dr. Santh Rani Thakur

Address: Division of Pharmacology, Institute of Pharmaceutical Technology, Sri Padmavathi Mahila Visvavidyalayam, Tirupati- 517502, Andhra Pradesh, India

Email ✉: drsanthrani@gmail.com

Tel.: +91-9849077507

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Sathuluri Vineela *et al.* This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Assessment of *in - vitro* Anti-inflammatory Activity

Inhibition of Albumin Denaturation

The anti-inflammatory activity of SA was studied by using inhibition of albumin denaturation method according to Mizushima and Kobayashi M., 1968; Sakat et al. 2010^[19,20] with minor modifications. The reaction mixture consisted of SA in different concentrations and 1% aqueous solution of bovine albumin fraction, pH of the reaction mixture was adjusted using 1N HCl. The samples were incubated at 37°C for 20 minutes and then heated to 51°C for 20 minutes, after cooling, the sample's turbidity was measured at 660 nm. The experiment was performed in triplicate. The percentage inhibition of protein denaturation was calculated using the following formula:

$$\text{Percentage inhibition} = (\text{Absorbance of Control} - \text{Absorbance of Sample}) / \text{Absorbance of control} \times 100$$

Membrane stabilization

Preparation of Red Blood Cell (RBCs) Suspension

Sheep blood was collected from the slaughter house 1 hour before the experiment and transferred to centrifuge tubes. The tubes were centrifuged at 3000 rpm for 10 min and were washed three times with equal volume of normal saline. The volume of blood was reconstituted as 10 % v/v suspension with normal saline.^[19]

Heat-induced Haemolysis

The reaction mixture (2 mL) consisted of 1 mL SA in different concentrations (10-500 µg/mL) and 1 mL of 10% RBCs suspension. Instead of SA, only saline was added to the control test tube. Aspirin (200 µg/mL) was used as a standard drug. Centrifuge tubes containing reaction mixture were incubated in a water bath at 56°C for 30 minutes. At the end of incubation, the tubes were cooled under running tap water. The reaction mixture was centrifuged at 2500 rpm for 5 minutes, and the absorbance of the supernatants was measured at 560 nm.^[19,20] The experiment was performed in triplicate for all the test samples. The Percentage inhibition of hemolysis was calculated as follows:

$$\text{Percentage inhibition} = (\text{Absorbance of Control} - \text{Absorbance of Sample}) / \text{Absorbance of Control} \times 100$$

Hypotonicity-induced Hemolysis

Different SA concentrations, standard sample and control were separately mixed with 1ml phosphate buffer, 2 mL of hypersaline, and 0.5 mL of HRBC suspension. Aspirin (200 µg/ml) was used as a standard drug. Assay mixtures were incubated at 37°C for 30 minutes and centrifuged at 3000 rpm. The supernatant liquid was decanted, and a Spectrophotometer estimated the hemoglobin content at 560 nm.^[19,20] The percentage hemolysis was estimated by assuming the hemolysis produced in the control as 100%.

$$\text{Percentage protection} = 100 - (\text{OD sample} / \text{OD control}) \times 100$$

Assay of Cyclooxygenase-2 and 5-Lipoxygenase Inhibition

Lymphocyte culture preparation

Human peripheral lymphocytes were cultured in RPMI 1640 [HIMEDIA] media, supplemented with 20% heat-inactivated fetal bovine serum, antibiotics (Penicillin and Streptomycin). Phyto haemagglutinin (HIMEDIA) was used as the stimulant for cell proliferation. The culture was filtered using 0.2 µm pore sized cellulose acetate filter (Sartorius) in completely aseptic conditions. Fresh plasma was aseptically added to the culture at a concentration of 1 x 10⁶ cells/ml. The culture was then incubated for 72 hours. After 24 hours, culture is activated by adding 1 µL lipopolysaccharide and incubated for 24 hours. Ethyl acetate and aqueous extracts were added in a final concentration of 100, 500 µg/mL and incubated for 24 hours. Ibuprofen at a concentration of 100 µg/mL was added and incubated for 24 hours. After incubation, the cells were pelleted by centrifugation. The isolation was done by spinning at 6000 rpm for 10 minutes. The supernatant was discarded and 50 µL of cell lysis buffer was added and again centrifuged at 6000 rpm for 10 minutes. The supernatant was discarded and the anti-inflammatory assay was done using pellet suspended in required amount of supernatant.^[21]

Assay of Cyclooxygenase-2

The assay mixture contained Tris- HCl buffer, glutathione, hemoglobin and enzyme. The assay was initiated by the addition of 200 µM of arachidonic acid incubated at 37°C and terminated after 20 min by the addition of 0.2 mL of 10% trichloroacetic acid (TCA) in 1N HCl, mixed and 0.2 mL of thiobarbiturate (TBA) was added and contents heated in a boiling water bath for 20 minutes, cooled and centrifuged at 1000 rpm for 3 minutes. The supernatant was measured at 632 nm for COX activity.^[21]

Assay of 5-Lipoxygenase

70 mg of linoleic acid and equal weight of tween 20 was dissolved in 4 mL of oxygen-free water and mixed back and forth with a pipette avoiding air bubbles. A sufficient amount of 0.5 N sodium hydroxide was added to yield a clear solution and then made up to 25 mL using oxygen free water. This was divided into 0.5 mL portions and flushed with nitrogen gas before closing and kept frozen until needed. The reaction was carried out in a quartz cuvette at 25°C with 1-cm light path. The assay mixture had 2.75 mL Tris buffer of pH 7.4, 0.2 mL of sodium linoleate, and 50 µL of the enzyme. OD was measured in 234 nm.^[21]

Statistical Analysis

Data were expressed as mean ± SEM (n = 3) and analyzed using one way analysis of variance (ANOVA) followed by Dunnet's T test using Graph pad prism 8.0. A value of p < 0.05 was considered to be statistically significant.

RESULTS

Effect of Sinapic Acid on Albumin Denaturation:

Sinapic acid significantly ($p < 0.001$) inhibited the albumin denaturation in a dose-dependent manner and maximum inhibition (70 %) was observed at 500 $\mu\text{g/mL}$ as compared with control. Aspirin, the standard drug used in this study showed maximum albumin denaturation inhibition (68.42 %) at 200 $\mu\text{g/mL}$ when compared with control ($p < 0.001$) (Fig. 1).

Effect of Sinapic Acid on Heat-induced Hemolysis

Sinapic acid dose-dependently increased the percentage inhibition of heat-induced hemolysis. Sinapic acid significantly ($p < 0.001$) inhibited (57.14 %) the heat-induced hemolysis at 500 $\mu\text{g/mL}$ when compared with control. Aspirin also significantly ($p < 0.001$) inhibited (70.12 %) the heat-induced hemolysis at 200 $\mu\text{g/mL}$ when compared with control (Fig. 2).




Fig 1: Effect of Sinapic acid on albumin denaturation
Values were expressed as mean $\pm$ SEM ($n = 3$); *** ($P < 0.001$) Vs Control group




Fig 2: Effect of Sinapic acid on heat induced haemolysis
Values were expressed as mean $\pm$ SEM ($n = 3$); *** ($p < 0.001$) Vs Control group

Effect of Sinapic Acid on Hypotonicity induced Hemolysis

Sinapic acid significantly ($p < 0.001$) protected hypotonic solution induced lysis of the erythrocyte membrane in a dose-dependent manner and maximum inhibition of hemolysis (82.06 %) was observed at 500 $\mu\text{g/mL}$ as compared with control. Aspirin at 200 $\mu\text{g/mL}$ showed significantly ($p < 0.001$) protection (70.12%) against the damaging effect of a hypotonic solution when compared with control (Fig. 3).

Effect of Sinapic Acid on Cyclooxygenase-2

Sinapic acid significantly ($p < 0.001$) inhibited Cyclooxygenase-2 and maximum inhibition of Cyclooxygenase-2 (71.77 %) was observed at 500 $\mu\text{g/mL}$ as compared with control. Ibuprofen at 100 $\mu\text{g/mL}$ showed significant ($P < 0.001$) inhibition of Cyclooxygenase-2 (87.88 %) (Fig. 4).




Fig 3: Effect of Sinapic acid on hypotonicity induced haemolysis
Values were expressed as mean $\pm$ SEM ($n = 3$); *** ($p < 0.001$) Vs Control group




Fig 4: Effect of Sinapic acid on Cyclooxygenase-2
Values were expressed as Mean $\pm$ SEM ($n = 3$); ** ($p < 0.01$), *** ($p < 0.001$) Vs Std group






Fig 5: Effect of Sinapic acid on 5-lipoxygenase
Values were expressed as mean $\pm$ SEM (n = 3); ***($p < 0.001$) Vs Standard group

Effect of Sinapic Acid on 5-lipoxygenase

Sinapic acid significantly ($p < 0.001$) inhibited lipoxygenase and maximum inhibition of 5-lipoxygenase (77.66 %) was observed at 500 $\mu\text{g/mL}$ as compared with control. Ibuprofen at 100 $\mu\text{g/mL}$ showed significant ($p < 0.001$) inhibition of lipoxygenase (70.62 %) (Fig. 5).

DISCUSSION

Inflammation may be a pathophysiological state related to pain. The free nerve endings of peripheral nerve fibers in general tissues respond to inflammatory factors, like pH, bradykinin, histamine or prostaglandins, by generating electrical activity which is perceived as pain,^[22] this useful pain state helps to safeguard us from adverse environments and might be managed by over-the-counter medications.^[23] Arachidonic acid undergoes 2 metabolic pathways: the cyclooxygenase (COX) pathway involving cyclooxygenase-1 (COX-1) and cyclooxygenase-2 (COX-2) to synthesise the prostaglandins and thromboxanes^[24] and the lipoxygenase (LOX) pathway, involving 5-lipoxygenase (5-LOX), 12-lipoxygenase (12-LOX) and 15-lipoxygenase (15-LOX), to produce the leukotrienes and hydroperoxy fatty acids.^[25,26] COX and LOX pathways participate in the induction of numerous pathologies, particularly inflammatory diseases. These include arthritis,^[27] fever, chronic pain,^[28] sepsis, burn injury,^[29] inflammatory bowel disease,^[30] and malignant neoplastic disease processes in large intestine cancer.^[31] In recent years, the search for phytochemicals possessing anti-inflammatory properties is on rise because of their potential use within the medical care of assorted chronic and infectious diseases. In the present study, we investigated the anti-inflammatory potential of SA by *in-vitro* strategies.

Protein Denaturation is a process in which proteins lose their tertiary structure and secondary structure by application of external stress or compound, such as strong acid or base, a concentrated inorganic salt, an

organic solvent, or heat. Most biological proteins lose their biological function when denatured.^[32] Protein denaturation is well correlated with the inflammatory response occurrence and leads to various inflammatory diseases.^[33] Tissue injury during life might be referred to as denaturation of cells' protein constituents or intercellular substances.^[34] Hence, a substance's ability to inhibit the denaturation of protein signifies apparent potential for anti-inflammatory activity. Sinapic acid effectively inhibited heat-induced albumin denaturation, indicating its anti-inflammatory property.

Membrane stabilization is used as a method to study the *in-vitro* anti-inflammatory activity, the erythrocyte membrane is analogous to the lysosomal membrane^[35,36] and its stabilization implies that the SA may well stabilize lysosomal membranes. Stabilization of lysosomal membrane is important in limiting the inflammatory response by preventing the release of lysosomal constituents of activated neutrophils, such as bacterial enzymes and proteases, further aggravating tissue inflammation and damage.^[37] The lysosomal enzymes released during inflammation produce various disorders. The extracellular activity of these enzymes is related to acute or chronic inflammation. The non-steroidal drugs act either by inhibiting these lysosomal enzymes or by stabilizing the lysosomal membrane.^[38] SA significantly inhibited the heat and hypotonicity induced hemolysis.

Medications that inhibit both COX-1/2 and 5-LOX pathways are believed to be superior to conventional NSAIDs, they produce a synergistic effect and achieve optimal anti-inflammatory activity by blocking the production of both leukotrienes and prostaglandins.^[39,40] This is supported by a significant reduction of collagen-induced arthritis in animal models upon concomitant administration of NSAID and leukotriene synthesis inhibitors.^[40,41] 5-LOX is the first and the key enzyme involved in the arachidonic acid pathway to produce leukotrienes.^[42] The 5-LOX pathway is associated with various diseases, including asthma,^[43] inflammatory bowel disease,^[44] cancers (prostate, pancreatic, and breast);^[45,46] and cardiovascular diseases including atherosclerosis, heart attack and stroke.^[47,48] COX and LOX inhibitors are considered promising agents for the treatment of inflammatory diseases on the basis of the important roles of LOX pathways.^[49] In this study, SA significantly inhibited both COX and LOX, indicating its anti-inflammatory activity.

Steroidal drugs are more potent anti-inflammatory agents but have adverse metabolic effects; NSAIDs have a dose ceiling effect and are not devoid of numerous adverse effects. 5LOX and CoX1/2 inhibitors are more potent anti-inflammatory agents. Sinapic Acid was screened *in-vitro* for anti-inflammatory activity using standard protocols. The results are encouraging as it blocks LOX COX and protein denaturation. Our results indicate that sinapic acid

possesses good anti-inflammatory activity by inhibiting protein denaturation, hemolysis, Cyclooxygenase-2 and lipoxygenase, indicating its potential as novel lead molecule for inflammation. Hence, further *in-vivo* studies are required to explore Sinapic acid's mechanism as anti-inflammatory agent.

ACKNOWLEDGMENTS

The authors are grateful to the management of Chebrolu Hanumaiah Institute of Pharmaceutical Sciences for extending their support and providing facilities to complete this work.

REFERENCES

- Medzhitov R. Origin and physiological roles of inflammation. *Nature*. 2008 Jul;454(7203):428-35.
- Serhan CN. Resolution phase of inflammation: novel endogenous anti-inflammatory and proresolving lipid mediators and pathways. *Annu. Rev. Immunol.*. 2007 Apr 23;25:101-37.
- Schett G. Rheumatoid arthritis: inflammation and bone loss. *Wiener Medizinische Wochenschrift*. 2006 Jan 1;156(1-2):34-41.
- Karin M, Greten FR. NF- κ B: linking inflammation and immunity to cancer development and progression. *Nature reviews immunology*. 2005 Oct;5(10):749-59.
- Libby P, Ridker PM, Maseri A. Inflammation and atherosclerosis. *Circulation*. 2002 Mar 5;105(9):1135-43.
- Hickson RC, Ball KL, Falduto MT. Adverse effects of anabolic steroids. *Medical toxicology and adverse drug experience*. 1989 Aug 1;4(4):254-71.
- Wongrakpanich S, Wongrakpanich A, Melhado K, Rangaswami J. A comprehensive review of non-steroidal anti-inflammatory drug use in the elderly. *Aging and disease*. 2018 Feb;9(1):143.
- Sawa T, Nakao M, Akaike T, Ono K, Maeda H. Alkylperoxyl radical-scavenging activity of various flavonoids and other phenolic compounds: implications for the anti-tumor-promoter effect of vegetables. *Journal of Agricultural and Food Chemistry*. 1999 Feb 15;47(2):397-402.
- Kuwahara H, Kanazawa A, Wakamatsu D, Morimura S, Kida K, Akaike T, Maeda H. Antioxidative and antimutagenic activities of 4-vinyl-2, 6-dimethoxyphenol (canolol) isolated from canola oil. *Journal of agricultural and food chemistry*. 2004 Jul 14;52(14):4380-7.
- D. A. Moreno, S. P'erez-Balibrea, F. Ferreres, A. Gil-Izquierdo, and C. Garc'ia-Viguera. "Acylated anthocyanins in broccoli sprouts," *Food Chemistry*, 2010 vol. 123, no. 2, pp. 358–363.
- Koski A, Pekkarinen S, Hopia A, Wähälä K, Heinonen M. Processing of rapeseed oil: effects on sinapic acid derivative content and oxidative stability. *European Food Research and Technology*. 2003 Aug 1;217(2):110-4.
- Engels C, Schieber A, Gänzle MG. Sinapic acid derivatives in defatted Oriental mustard (*Brassica juncea* L.) seed meal extracts using UHPLC-DAD-ESI-MS n and identification of compounds with antibacterial activity. *European Food Research and Technology*. 2012 Mar 1;234(3):535-42.
- Maddox CE, Laur LM, Tian L. Antibacterial activity of phenolic compounds against the phytopathogen *Xylella fastidiosa*. *Current microbiology*. 2010 Jan 1;60(1):53.
- Kikuzaki H, Hisamoto M, Hirose K, Akiyama K, Taniguchi H. Antioxidant properties of ferulic acid and its related compounds. *Journal of agricultural and food chemistry*. 2002 Mar 27;50(7):2161-8.
- Hudson EA, Dinh PA, Kokubun T, Simmonds MS, Gescher A. Characterization of potentially chemopreventive phenols in extracts of brown rice that inhibit the growth of human breast and colon cancer cells. *Cancer Epidemiology and Prevention Biomarkers*. 2000 Nov 1;9(11):1163-70.
- Kanchana G, Shyni WJ, Rajadurai M, Periasamy R. Evaluation of antihyperglycemic effect of sinapic acid in normal and streptozotocin-induced diabetes in albino rats. *Global Journal of Pharmacology*. 2011;5(1):33-9.
- Sun XL, Ito H, Masuoka T, Kamei C, Hatano T. Effect of *Polygala tenuifolia* root extract on scopolamine-induced impairment of rat spatial cognition in an eight-arm radial maze task. *Biological and Pharmaceutical Bulletin*. 2007 Sep 1;30(9):1727-31.
- Yoon BH, Jung JW, Lee JJ, Cho YW, Jang CG, Jin C, Oh TH, Ryu JH. Anxiolytic-like effects of sinapic acid in mice. *Life sciences*. 2007 Jun 27;81(3):234-40.
- Mizushima Y, Kobayashi M. Interaction of anti-inflammatory drugs with serum proteins, especially with some biologically active proteins. *Journal of Pharmacy and Pharmacology*. 1968 Mar;20(3):169-73.
- Sakat S, Tupe P, Juvekar A. Gastroprotective effect of methanol extract of *Oxalis corniculata* Linn (whole plant) experimental animals. *Planta Medica*. 2010 Aug;76(12):P090.
- Viji V, Helen A. Inhibition of lipoxygenases and cyclooxygenase-2 enzymes by extracts isolated from *Bacopa monniera* (L.) Wettst. *Journal of ethnopharmacology*. 2008 Jul 23;118(2):305-11.
- Jang Y, Kim M, Hwang SW. Molecular mechanisms underlying the actions of arachidonic acid-derived prostaglandins on peripheral nociception. *Journal of Neuroinflammation*. 2020 Dec 1;17(1):30.
- José Oswaldo de Oliveira Júnior, Caio Sander Andrade Portella Junior, Cláudia Panossian Cohen. Inflammatory mediators of neuropathic pain, *Rev Dor. São Paulo*, 2016;17 (Suppl 1):S35-42.
- Rainsford KD. Anti-inflammatory drugs in the 21st century. In *Inflammation in the pathogenesis of chronic diseases 2007* (pp. 3-27). Springer, Dordrecht.
- Samuelsson B, Dahlen SE, Lindgren JA, Rouzer CA, Serhan CN. Leukotrienes and lipoxins: structures, biosynthesis, and biological effects. *Science*. 1987 Sep 4;237(4819):1171-6.
- Yedgar S, Krinsky M, Cohen Y, Flower RJ. Treatment of inflammatory diseases by selective eicosanoid inhibition: a double-edged sword?. *Trends in pharmacological sciences*. 2007 Sep 1;28(9):459-464.
- Honda T, Segi-Nishida E, Miyachi Y, Narumiya S. Prostacyclin-IP signaling and prostaglandin E2-EP2/EP4 signaling both mediate joint inflammation in mouse collagen-induced arthritis. *The Journal of experimental medicine*. 2006 Feb 20;203(2):325-35.
- Mabuchi T, Kojima H, Abe T, Takagi K, Sakurai M, Ohmiya Y, Uematsu S, Akira S, Watanabe K, Ito S. Membrane-associated prostaglandin E synthase-1 is required for neuropathic pain. *Neuroreport*. 2004 Jun 28;15(9):1395-8.
- Hahn EL, He LK, Gamelli RL. Prostaglandin E2 synthesis and metabolism in burn injury and trauma. *Journal of Trauma and Acute Care Surgery*. 2000 Dec 1;49(6):1147-54.
- Subbaramaiah K, Yoshimatsu K, Scherl E, Das KM, Glazier KD, Golijanin D, Soslow RA, Tanabe T, Naraba H, Dannenberg AJ. Microsomal prostaglandin E synthase-1 is overexpressed in inflammatory bowel disease Evidence for involvement of the transcription factor Egr-1. *Journal of Biological Chemistry*. 2004 Mar 26;279(13):12647-58.
- Backlund MG, Mann JR, DuBois RN. Mechanisms for the prevention of gastrointestinal cancer: the role of prostaglandin E2. *Oncology*. 2005;69(Suppl. 1):28-32.
- Mizushima Y. Screening test for antirheumatic drugs. *The Lancet*. 1966 Aug 20;288(7460):443.
- Sangeetha G, Vidhya R. In vitro anti-inflammatory activity of different parts of *Pedaliu murex* (L.). *inflammation*. 2016;4:5.
- Opie EL. On the relation of necrosis and inflammation to denaturation of proteins. *The Journal of Experimental Medicine*. 1962 Feb 28;115(3):597.
- Gandhidasan R, Thamaraichelvan A, Baburaj S. Anti inflammatory action of *Lannea coromandelica* by HRBC membrane stabilization. *Fitoterapia*. 1991;62(1):81-3.
- Shenoy S, Shwetha K, Prabhu K, Maradi R, Bairy KL, Shanbhag T. Evaluation of anti-inflammatory activity of *Tephrosia purpurea*



- in rats. Asian Pacific journal of tropical medicine. 2010 Mar 1;3(3):193-5.
37. Leelaprakash, G. and Dass, S.M.. Invitro anti-inflammatory activity of methanol extract of *Enicostemma axillare*. *International Journal of Drug Development and Research*, 2011 3(3), pp.189-196.
 38. Vadivu R, Lakshmi KS. In vitro and in vivo anti-inflammatory activity of leaves of *Symplocos cochinchinensis* (Lour) Moore ssp Laurina. ||| Bangladesh Journal of Pharmacology|||. 2008 Aug 1;3(2):121-4.
 39. Celotti F, Laufer S. Anti-inflammatory drugs: new multitarget compounds to face an old problem. The dual inhibition concept. *Pharmacological Research*. 2001 May 1;43(5):429-36.
 40. Martel-Pelletier J, Lajeunesse D, Reboul P, Pelletier JP. Therapeutic role of dual inhibitors of 5-LOX and COX, selective and non-selective non-steroidal anti-inflammatory drugs. *Annals of the rheumatic diseases*. 2003 Jun 1;62(6):501-9.
 41. Nickerson-Nutter CL, Medvedeff ED. The effect of leukotriene synthesis inhibitors in models of acute and chronic inflammation. *Arthritis & Rheumatism: Official Journal of the American College of Rheumatology*. 1996 Mar;39(3):515-21.
 42. Zhang YY, Walker JL, Huang A, KEANEY Jr JF, Clish CB, Serhan CN, Loscalzo J. Expression of 5-lipoxygenase in pulmonary artery endothelial cells. *Biochemical Journal*. 2002 Jan 15;361(2):267-276.
 43. Charlier C, Michaux C. Dual inhibition of cyclooxygenase-2 (COX-2) and 5-lipoxygenase (5-LOX) as a new strategy to provide safer non-steroidal anti-inflammatory drugs. *European journal of medicinal chemistry*. 2003 Jul 1;38(7-8):645-59.
 44. Jupp J, Hillier K, Elliott DH, Fine DR, Bateman AC, Johnson PA, Cazaly AM, Penrose JF, Sampson AP. Colonic expression of leukotriene-pathway enzymes in inflammatory bowel diseases. *Inflammatory bowel diseases*. 2007 May 1;13(5):537-46.
 45. Avis I, Hong SH, Martínez A, Moody T, Choi YH, Trepel J, Das R, Jett M, Mulshine JL. Five-lipoxygenase inhibitors can mediate apoptosis in human breast cancer cell lines through complex eicosanoid interactions. *The FASEB Journal*. 2001 Sep;15(11):2007-9.
 46. Romano M, Catalano A, Nutini M, D'URBANO ET, Crescenzi C, Claria J, Libner R, Davi G, Procopio A. 5-Lipoxygenase regulates malignant mesothelial cell survival: involvement of vascular endothelial growth factor. *The FASEB Journal*. 2001 Nov;15(13):2326-36.
 47. Funk CD. Leukotriene modifiers as potential therapeutics for cardiovascular disease. *Nature Reviews Drug Discovery*. 2005 Aug;4(8):664-72.
 48. Mehrabian M, Allayee H. 5-lipoxygenase and atherosclerosis. *Curr Opin Lipidol* 2003;14: 447–457.
 49. Naveau B. Dual inhibition of cyclo-oxygenases and 5-lipoxygenase: a novel therapeutic approach to inflammation?. *Joint, bone, spine: revue du rhumatisme*. 2005 May; 72(3):199.

HOW TO CITE THIS ARTICLE: Vineela S, Devi CM, Thakur SR. Evaluation of *in-vitro* anti-inflammatory activity of sinapic acid. *Int. J. Pharm. Sci. Drug Res.* 2021;13(1):30-35. **DOI:** 10.25004/IJPSDR.2021.130105