

## Review on Effects of Organophosphate Pesticides in Human Health and Way to Mitigates its Harmful Impacts



Mr. Subrata Kumar Payra<sup>1</sup>, Ms. Deepa Goswami<sup>2</sup>, Mrs. Sanchita Pan<sup>3</sup>, Mr. Atanu Bera<sup>4</sup>,  
Ms. Supriya Jana<sup>5</sup>, Mr. Jagmohan Sing<sup>6</sup>, Mr. Atanu Bhowmik<sup>7</sup>

<sup>1</sup>Assistant Professor, Zoology Department, Panskura Banamali College, 721152, West Bengal, India.

<sup>2</sup>PhD Scholar, Chemical Engineering Department, Jadavpur University, Kolkata, 700032.

<sup>3</sup>SACT, Zoology Department, Jhargram Raj College, Jhargram, 721507.

<sup>4</sup>M.Sc, PhD Scholar under Vidyasagar University, Medinipur, 721102.

<sup>5,6</sup>B.Sc. in Zoology, Jhargram Raj College, Jhargram, 721507.

<sup>7</sup>M.Sc, in Botany, From Vidyasagar University, Medinipur, 721102.

**ABSTRACT:** Due to increasing the human population day by day, for the control of the human population, needs to increase the agricultural productivity, for agricultural production is increasing we use pesticide. Pesticides not only kill the target pest species also affect non-target organisms. Pesticide pollution has become a significant problem all over the world. This is mainly due to the abuse and overuse of pesticides which remain in the soil for several years and leach into water bodies. Pesticides are chemical (synthetic or semi-synthetic) agents that control and destroy pest population. Organophosphate pesticides are major bulk that includes malathion, parathion, diazinon, fenthion, chlordecone, Imidacloprid, dichlorvos, chlorpyrifos, ethion, nerve gases (soman, sarin, tabun, VX), ophthalmic agents (echothiophate, isoflurophate), and antihelmintics (trichlorfon). Herbicides (tribufos or DEF, merphos) are tricresyl phosphate etc. Though various pesticides were banned due to its harmful impacts on human health and environment, but still some has been used in paddy fields, greenhouse culture, horticulture and even hydroponics. They persistent in nature (bio-accumulation and bio-magnification by the food chain, especially through pieces) and cause acute or chronic toxicity. They contain very noxious components that harm our neuronal system by cholinergic effects. Pesticide also has an unfavorable effect on aquatic organisms such as ataxia causing late maturation in fish. Various physical and chemical methods are used for the reduction the toxicity of pesticides, physically and chemically pesticide reduction process is costly and hazardous. One good strategy is applied for the remediation of pesticides is bioremediation, i.e- use of various modified strain of agricultural microbes especially bacteria.

**KEYWORDS:** Pesticides, Chlorpyrifos, Bio-magnification, Cholinergic effect and Bioremediation.

### INTRODUCTION

Organophosphates (OPs) are large group about 40% of total pesticide, (Rao et al., 2005) of chemical that used over past 7 decades in livestock practice, human health and welfare, and defensive agents. These are class of insecticides, several of which are highly toxic as contain cholinesterase inhibitor property thus directly or indirectly deals with CNS, i.e. Neurological disorders, (Gupta, 2001). Besides of this the direct exposure of it causes skin etching, seizure, respiratory disorders and ultimately death of organisms. The route of explosion may be differ in different organism, such as dermal adsorption, inhalation and through diet via food web in ecosystem (Costa, 2018). Thirty-six of OPs are presently registered for use in the United States, and all can potentially cause acute and sub-acute toxicity. Organophosphates are used in agricultural fields, homes, gardens and horticulture, and veterinary practices. Parathion and chlorpyrifos, has no longer registered for any use. All share a common mechanism of cholinesterase inhibition and can cause similar symptoms, although there are some differences within the class. However, that there is a wide range of toxicity in these agents and wide variation in dermal absorption. Organophosphates poison insects and other animals, including birds, amphibians and mammals (via ecological food chain), primarily by phosphorylation of the acetylcholinesterase enzyme (AChE) at nerve end plate. The result is a loss of available AChE so that the effectors organ becomes over- stimulated by the excess acetylcholine (ACh, the impulse-transmitting substance) in the nerve ending. At skeletal muscle junctions, excess ACh may be excitatory (cause muscle twitching) but may also weaken or paralyze the cell by depolarizing the end plate. Impairment of

## **Review on Effects of Organophosphate Pesticides in Human Health and Way to Mitigates its Harmful Impacts**

the diaphragm and thoracic skeletal muscles can cause respiratory paralysis (Routt and Roberts, 1999). In the CNS, high Ach concentrations cause sensory and behavioral disturbances, in coordination, depressed motor function and respiratory depression. Increased pulmonary secretions coupled with respiratory failure are the usual causes of death from organophosphate poisoning. Some of the most commonly reported early symptoms include headache, nausea, dizziness and hyper-secretion, the latter of which is manifested by sweating, salivation, lacrimation and rhinorrhea. Muscle twitching, weakness, tremor, in coordination, vomiting, abdominal cramps and diarrhea all signal worsening of the poisoned state. Organophosphate herbicides interacting with soil biota and altering the biomass of environment (Usman et al., 2017). Some organophosphates, such as diazinon, fenthion and methyl parathion, have significant lipid solubility, allowing fat storage with delayed toxicity due to late release.3,4 Delayed toxicity may also occur atypically with other organophosphates, specifically dichlorofenthion and demton-methy. Many organo-thiophosphates readily undergo conversion from thions (P=S) to oxons (P=O). Conversion occurs in the environment under the influence of oxygen and light and, in the body, chiefly by the action of liver microsomal enzymes. Level of threat are detected in various way: dose determination by LC50 and LD50 of OPs pesticide, various physio-chemical and biological parameter test of target tissue and symptoms (Zhang et al, 2017). Various degradation strategies should be taking now days like abiotic degradation by chemical oxidation, catalytic hydrolysis, adsorbing agents and gamma radiation. Major disadvantages of it are high cost and natural unsustainability. Another most effective and eco-friendly method is bioremediation by microbial strains on which more research and technology involved. In some cases the patient with severe exposure of OPs are treated with atropine like drugs. India is fourth largest producer of pesticide in the world that largely affects on India's GDP and economy. Pesticide Management Bill (PMB, 2020) is an opportunity to clean up India's food and farming system.

### **RESULT**

#### **1.1. The Impacts of Organophosphate Pesticides on Human Health:**

##### **Choline Esterase Effects:**

The cholinergic i.e. blocked the substrate binding site of Acetyl choline esterase enzyme, effects brought about by repeated administration of less than a single fatal dose are similar in type to the acute single-dose effects (WHO, 1986).

##### **Dermal or Cutaneous Effects:**

Shoot particle and pesticide droplets during spray at agricultural field directly expose at dermal contact, cause skin rashes and allergic manifestation.

##### **Immediate or Acute Toxicity:**

Acute toxicity is the ability of a substance to cause harmful impacts immediately in a single exposure or dose or any severe poisonous effect resulting from a single short-term exposure to a toxic substance. LD50 (lethal dose 50) is defined as the dose that kills 50% of a test population of the exposed animals (Ghosh and Philip, 2006). In an acute study, neurotoxic signs were seen at 190 mg/kg in rats, and were more prominent in females. Also, in vitro studies have shown butyl cholinesterase to be more sensitive to profenofos inhibition than acetyl cholinesterase (AChE) (McDaniel and Moser, 2004).

##### **Chronic Toxicity**

Chronic toxicity is the capacity of a substance to cause long-term or delayed adverse health effects. Several reports proved the chronic toxicity of atrazine on various test organisms. After the sign of respiratory distress and paralysis of the limbs, 40% rats were died at an atrazine oral dose of 20 mg/kg/day for 6 months. Body organs like brain, liver, kidneys, ovaries etc., were affected by organophosphates poisons and made structural and chemical changes, as well as growth retardation. In a study carried out for two years with dogs, results showed that 7.5 mg kg<sup>-1</sup> day<sup>-1</sup> atrazine dose caused decreasing in food intake and increased heart and liver weights. The NOEL on rat was 70 mg kg<sup>-1</sup> whereas in dog it was 15 mg kg<sup>-1</sup> of body weight (Sameeh, 2004).

##### **Long term Neuropathic Effects**

Delayed neuropathy has occurred occasionally in human being, livestock and experimental animals after intoxication with a variety of OPs esters. However, many OPP that might, theoretically, cause neuropathy, would only do so at a dose far above the lethal dose (WHO, 1986).

##### **Carcinogenic Effects**

Many OPs pesticides have not shown carcinogenic potential in animal experiments, but some pesticides do, through induction of tumors in rats and mice. No generalizations can be made due to some compounds exhibit mutagenic activity, whereas other compounds do not (WHO, 1986).

## Review on Effects of Organophosphate Pesticides in Human Health and Way to Mitigates its Harmful Impacts

### Reproductive Effects

A number of pesticides clearly have the potential to cause reproductive toxicity in animals, and several compounds e.g. (chlordecone) are known to affect human reproduction (Sameeh, 2004). Also OPs included insecticides (malathion, parathion, diazinon, fenthion, dichlorvos, chlorpyrifos, ethion), nerve gases (soman, sarin, tabun, VX), ophthalmic agents (echothiophate, isoflurophate), and antihelminthics (trichlorfon). Herbicides (tribufos [DEF], merphos) are tricresyl phosphate-containing industrial chemicals.

### Immunotoxicity

The scientific evidence suggesting that many pesticides damage the immune system. Animal studies have found that pesticides alter the immune system's normal structure, disturb immune responses, and reduce animal's resistance to antigens and infectious agents. In case of Malathion which is considered a very low toxic compound (oral LD50= 2100 mg/kg bw), for example, not regulates the immune system, especially affecting non-specific immune mechanisms (Sameeh, 2004).

### Cytogenetic Effects

Cytogenetic damage related to pesticides exposure has been reported in various populations. Some investigators have reported significant differences in the percentage of chromosomal aberrations (CAs) in exposed individuals (range, 2.66–10.30%) compared with control (range, 0.53– 5.52%) (Sameeh, 2004).

### Teratogenic Effects

Detailed data on the effects of organophosphate occupational exposure on pregnant women and their fetuses are not available, although such information would be valuable. In humans only a few cases of acute organophosphorus insecticide poisoning during pregnancy have been described. A 24-year-old woman in her third month of pregnancy injected herself with Malathion in a suicide attempt. A therapeutic abortion was performed 2 months later. Continuation of the pregnancy was considered to be dangerous, although the condition of the fetus was not described (WHO, 1986).

### Effects on the Immune System

The most Organophosphorus pesticides elicit autoimmune reactions and suppress the production of antibodies against various vaccines (Zackov, 1983).

### 1.2. Symptoms of Organophosphate Pesticide Poisoning (Acute to Chronic)

Level of Exposion	Mild Exposion	Moderate Exposion	High Exposion	Very High Exposion
Symptoms	Poor vision	Muscle Tremor	Slow heartbeat	Paralysis
	Nausea/Vomiting	Disorientation and balance	Coma	Fertility Disorder
	Muscle twitching	Diarrhoea	Narrow pupil	Cancer
	Abnormal Salivation	Cough	Excessive saliva, tears, and urination	Pancreatic disorders
	Headache	Sneezing	Neurological impaired	Digestive problems
	Weakness	Severe vomiting	Heart and lung abnormalities	Blood sugar
	Agitation		Difficulty in Breathing	Neurological disorder

Table. Shows level of exposure combine with manifested symptoms. Source: Article, A treatise on Organophosphate Pesticide pollution by Kaushal J. et al., 2020.

### 4.3 Bioremediation:

Bioremediation reduces pesticide contamination of agricultural soils by biodegradation processes via the metabolic activities of microorganisms. It is an efficient, cost-effective, and environment-friendly treatment. During the bioremediation processes, the microorganisms use the pesticides as co-substrates in their metabolic reactions together with other nutrients, thus eliminating them from the environment. The efficiency of these processes depends on the characteristics of pesticides, such as their distribution, their bioavailability, and their persistence in soil. It is necessary to promote the availability of pesticides to microorganisms: this is negatively affected by the adhesion of pesticides to soil particles and their low water solubility [Ortiz-

## Review on Effects of Organophosphate Pesticides in Human Health and Way to Mitigates its Harmful Impacts

Hernández et al, 2014]. In addition, the soil characteristics and the environmental conditions, such as pH, water content, microbial diversity, and temperature, influence the bioremediation efficacy.

Sl. No.	Organophosphate Pesticide	Microbes/ Microbial strain that metabolically degrade
1	Chlorpyrifos	<i>Bacillus pumilus</i> strain C2A1, <i>Pseudomonas aeruginosa</i> , <i>P. stutzeri</i> , <i>P. putida</i> , <i>Bacillus aryabhata</i> , <i>Stenotrophomonas</i> sp., <i>Leuconostoc mesenteroides</i> , <i>Trichoderma harzianum</i> , <i>Penicillium vermiculatum</i> , and <i>Mucor</i> sp. etc
2	Malathion	<i>Pseudomonas</i> sp., <i>P. stutzeri</i> , <i>Arthrobacter</i> , <i>Stenotrophomonas</i> sp., <i>Leuconostoc mesenteroides</i>
3	Diazinon	<i>Serratia marcescens</i> D1101. and <i>Pseudomonas</i> sp., <i>LactoBacillus brevis</i> , <i>Mycobacterium</i> sp., and <i>Stenotrophomonas</i> sp.
4	Methyl-parathion	<i>Pseudomonas</i> sp., <i>Xanthomonas</i> sp., <i>Pseudomonas</i> sp., <i>P. stutzeri</i> , <i>Arthrobacter</i> , <i>Stenotrophomonas</i> sp., <i>Leuconostoc mesenteroides</i> , <i>Brevis</i> L., <i>Plantarum</i> L., <i>Sakei</i> .
5	Fenitrothion	<i>Burkholderia</i> sp. strain NF1000, <i>Arthrobacter aurescens</i> TW17, <i>Corynebacterium</i> and <i>Arthrobacter</i> sp..
6	3-methyl-4- nitrophenol by	<i>Ralstonia</i> sp. SJ98

### CONCLUSION

Ever bursting human population demands more food, cloth and daily needs thus we must increase our livestock production many more in a poor illegal methods. That leads to the damage of microbial community and natural stability and sustainability, thus human health. Various controlling measurement should take but bioremediation or phytoremediation is most effective one. With modern researches, advance technologies, legislative strategy and human awareness we can protest against it. Controlling human population is one of most effective indirect method. If that is gone continuously the future generation will suffer most hazardously.

### ACKNOWLEDGEMENTS

The first author is thanks to his parents and beloved brother for supporting various purposes. The second author is grateful to her parents for continuous encouragement and supports. The third author is thanks to her teacher Debasish Chanda for his ideal teaching and also guiding to her. The fourth author owe his deepest sense of gratitude and regard to his teacher Dr. Pritiranjana Pahari, Assistant professor Co- coordinator, PG Department of Zoology, Tamralipta Mahavidyalaya for his suggestion, encouragement and support. The fifth author is thanks to her teacher Biswajit Ghosh, Assistant teacher in physics, Kharbandhi S.C. High School, Kharbandhi, W.B. for his ideal teaching and also guiding. The seventh author is thanks to his parent for continuous support.

**Conflict of Interest:** The Authors declare no conflict of interest.

**Funding:** The work doesn't require any external funding.

### REFERENCES

- 1) Abhijith BD, Ramesh M, Poopal RK (2016) Responses of metabolic and antioxidant enzymatic activities in gill, liver and plasma of *Catla catla* during methyl parathion exposure. *J Basic Appl Zool* 77:31–40.
- 2) Abhilash PC, Singh N (2009) Pesticide use and application: an Indian scenario. *J Hazard Mater* 165:1–12.
- 3) Abraham J, Silambarasan S (2016) Biodegradation of chlorpyrifos and its hydrolysis product 3,5,6-trichloro-2-pyridinol using a novel bacterium *Ochrobactrum* sp. *JAS2: a proposal of its metabolic pathway. Pestic Biochem Physiol* 126:13–21.

## Review on Effects of Organophosphate Pesticides in Human Health and Way to Mitigates its Harmful Impacts

- 4) Akbar S, Sultan S (2016) Soil bacteria showing a potential of chlorpyrifos degradation and plant growth enhancement. *Braz J Microbiol* 47(3):563–570.
- 5) Akbar S, Sultan S, Kertesz M (2014) Bacterial community analysis in chlorpyrifos enrichment cultures via DGGE and use of bacterial consortium for CP biodegradation. *World J Microbiol Biotechnol* 30:2755–2766.
- 6) Alavanja MC, Samanic C, Dosemeci M, Lubin J, Tarone R, Lynch CF, Knott C, Thomas K, Hoppin JA, Barker J, Coble J, Sandler DP, Blair A (2003) Use of agricultural pesticides and prostate cancer risk in the agricultural health study cohort. *Am J Epidemiol* 157:800–814.
- 7) Anwar S, Liaquat F, Khan QM, Khalid ZM, Iqbal S (2009) Biodegradation of chlorpyrifos and its hydrolysis product 3,5,6-trichloro-2-pyridinol by *Bacillus pumilus* strain C2A1. *J Hazard Mater* 168(1):400–405.
- 8) Barcelo D (1991) Occurrence, handling and chromatographic determination of pesticides in the aquatic environment. A review. *Analyst* 116(7):681–689.
- 9) Bharagava RN, Chowdhary P, Saxena G (2017a) Bioremediation: an ecosustainable green technology: its applications and limitations. In: Bharagava RN (ed) *Environmental pollutants and their bioremediation approaches*, 1st edn. CRC Press/Taylor & Francis Group, Boca Raton, pp 1–22. <https://doi.org/10.1201/9781315173351-2>.
- 10) Blakley BR, Yole MJ, Brousseau P, Boermans H, Fournier M (1999) Effect of chlorpyrifos on immune function in rats. *Vet Hum Toxicol* 41(3):140–144.
- 11) Chandra R, Saxena G, Kumar V (2015) Phytoremediation of environmental pollutants: an eco-sustainable green technology to environmental management. In: Chandra R (ed) *Advances in biodegradation and bioremediation of industrial waste*, 1<sup>st</sup> edn. CRC Press/Taylor & Francis Group, Boca Raton, pp 1–30. <https://doi.org/10.1201/b18218-2>.
- 12) Charoensri K, Esuchart U, Nouwarath S, Pairote P (2001) Degradation of methyl parathion in an aqueous medium by soil bacteria. *Sci Asia* 27:261–271.
- 13) Chaudhry GR, Ali AN, Wheeler WB (1988) Isolation of a methyl parathion-degrading *Pseudomonas* sp. that possesses DNA homologous to the opd gene from a *Flavobacterium* sp. *Appl Environ Microbiol* 54:288–293
- 14) Deng S, Chen Y, Wang D, Shi T, Wu X, Ma X, Li X, Hua R, Tang X, Li QX (2015) Rapid biodegradation of organophosphorus pesticides by *Stenotrophomonas* sp. G1. *J Hazard Mater* 297:17–24.
- 15) Dhanjal NIK, Kaur P, Sud D, Cameotra SS (2014) Persistence and biodegradation of quinalphos using soil microbes. *Water Environ Res* 86:457–461.
- 16) Elersek T, Filipic M (2011) Organophosphorous pesticides – mechanisms of their toxicity. In: Stoytcheva M (ed) *Pesticides – the impacts of pesticides exposure*.
- 17) Engel LS, Hill DA, Hoppin JA, Lubin JH, Lynch CF, Pierce J, Samanic C, Sandler DP, Blair A, Alavanja MC (2005) Pesticide use and breast cancer risk among farmers' wives in the agricultural health study. *Am J Epidemiol* 161:121–135.
- 18) EPA (2020) Environmental protection agency reregistration eligibility decision for profenofos. [http://www.epa.gov/oppsrrd1/REDs/profenofos\\_red.pdf](http://www.epa.gov/oppsrrd1/REDs/profenofos_red.pdf).
- 19) Fawzy I, Iman Z, Hamza A (2007) The effect of an Organophosphorus insecticide on the hepatic, renal and pulmonary tissues of mice fetuses Egypt. *J Med Lab Sci* 16:99–113.
- 20) Feng F, Ge J, Li Y, Cheng J, Zhong J, Yu X (2017) Isolation, colonization, and Chlorpyrifos degradation mediation of the endophytic bacterium *Sphingomonas* strain HJY in Chinese chives (*Allium tuberosum*). *J Agric Food Chem* 65(6):1131–1138.
- 21) Gangireddygar VSR, Kalva PK, Ntushelo K, Bangeppagari M, Djami Tchatchou A, Bontha RR (2017) Influence of environmental factors on biodegradation of quinalphos by *Bacillus thuringiensis*. *Environ Sci Eur* 29(1):11.
- 22) Ghanem I, Orfi M, Shamma M (2007) Biodegradation of chlorpyrifos by *Klebsiella* sp. isolated from an activated sludge sample of waste water treatment plant in Damascus. *Folia Microbiol* 52(4):423–427.
- 23) Gilani RA, Rafique M, Rehman A, Munis MFH, ur Rehman S, Chaudhary HJ (2016) Biodegradation of chlorpyrifos by bacterial genus *Pseudomonas*. *J Basic Microbiol* 56:105–119.
- 24) Gupta R. C., 2001, *Toxicology of Organophosphate and Carbamate compounds*, pp. 5-14.
- 25) Huang QY, Huang L, Huang HQ (2011) Proteomic analysis of methyl parathion-responsive proteins in zebrafish (*Danio rerio*) brain. *Comp Biochem Physiol C Toxicol Pharmacol* 153(1):67–74.
- 26) Ishag AESA, Abdelbagi AO, Hammad AMA, Elsheikh EAE, Elsaid OE, Hur J-H, Laing MD (2016) Biodegradation of chlorpyrifos, malathion and dimethoate by three strains of bacteria isolated from pesticide-polluted soils in Sudan. *J Agric Food Chem* 64:8491–8498.
- 27) John EM, Sreekumar J, Jisha MS (2016) Optimization of Chlorpyrifos degradation by assembled bacterial consortium using response surface methodology. *Soil Sedimentol Contam* 25:668–682.

## Review on Effects of Organophosphate Pesticides in Human Health and Way to Mitigates its Harmful Impacts

- 28) Malghani S, Chatterjee N, Hu X, Zejiao L (2009a) Isolation and characterization of a profenofos degrading bacterium. *J Environ Sci (China)* 21:1591–1597.
- 29) M. A. Ghorab, M. S. Khalil, 2015, Toxicological Effects of Organophosphates Pesticides. doi: 10.11648/j.ijema.20150304.13. *International Journal of Environmental Monitoring and Analysis*.
- 30) Mohapatra PK (2008) Textbook of environmental microbiology. I.K. International Publishing House Pvt. Ltd, New Delhi.
- 31) Nair AM, Rebello S, Rishad KS, Asok AK, Jisha MS (2015) Biosurfactant facilitated biodegradation of quinalphos at high concentrations by *Pseudomonas aeruginosa* Q10. *Soil Sediment Contam* 24:542–553.
- 32) Ortiz-Hernández, M.L.; Rodríguez, A.; Sánchez-Salinas, E.; Castrejón-Godínez, M.L. Bioremediation of Soils Contaminated with Pesticides: Experiences in Mexico. In *Bioremediation in Latin America: Current Research and Perspectives*; Alvarez, A., Polti, M.A., Eds.; Springer: Cham, Switzerland, 2014; ISBN 9783319057385
- 33) Prakash A, Khan S, Aggarwal M, Telang AG, Malik JK (2009) Chlorpyrifos induces apoptosis in murine thymocytes. *Toxicol Lett* 189:S83.
- 34) Rani NL, Lalitha-kumari D (1994) Degradation of methyl parathion by *Pseudomonas putida*. *Can J Microbiol* 4:1000–1004.
- 35) Salunkhe VP, Sawant IS, Banerjee K, Rajguru YR, Wadkar PN, Oulkar DP, Naik DG, Sawant SD (2013) Biodegradation of profenofos by *Bacillus subtilis* isolated from grapevines (*Vitis vinifera*). *J Agric Food Chem* 61:7195–7202.
- 36) Singh BK, Walker A, Morgan JAW, Wright DJ (2003) Effect of soil pH on the biodegradation of chlorpyrifos and isolation of a chlorpyrifos-degrading bacterium. *Appl Environ Microbiol* 69:5198–5206.
- 37) Suleiman U., Kundiri A. M., maximilien N.. 2017. Effects of Organophosphate Herbicides on Biological Organisms in Soil Medium, a mini review, *J. Ecol. Toxicol.*, 1 (2017).
- 38) Tallur PN, Mulla SI, Megadi VB, Talwar MP, Ninnekar HZ (2015) Biodegradation of cypermethrin by immobilized cells of *Micrococcus* sp. strain CPN 1. *Braz J Microbiol* 46(3):667–672.
- 39) Tilman D, Cassman KG, Matson PA, Naylor R, Polasky S (2002) Agricultural sustainability and intensive production practices. *Nature* 418:671–677.
- 40) Venkateswarl S., C. H., Eddleston M., 2005 Pesticide Poisoning in South India: Opportunities for Prevention and Improved Medical Management, *Trop. Med. Int. Health*, 10 (2005), pp. 581-588.
- 41) WHO (2004) Methyl parathion in drinking-water. WHO/SDE/WSH/03.04/106. [http://www.who.int/water\\_sanitation\\_health/dwq/chemicals/methylparathion.pdf](http://www.who.int/water_sanitation_health/dwq/chemicals/methylparathion.pdf).
- 42) Yadav M, Shukla AK, Srivastva N, Upadhyay SN, Dubey SK (2016) Utilization of microbial community potential for removal of chlorpyrifos: a review. *Crit Rev Biotechnol* 36(4):727–742.
- 43) Zheng Y, Long L, Fan Y, Gan J, Fang J, Jin W (2013) A review on the detoxification of organophosphorus compounds by microorganisms. *Afr J Microbiol Res* 7:2127–2134.
- 44) Zhongli C, Shunpeng L, Guoping F (2001) Isolation of methyl parathion-degrading strain M6 and cloning of the methyl parathion hydrolase gene. *Appl Environ Microbiol* 67:4922–4925.



There is an Open Access article, distributed under the term of the Creative Commons Attribution – Non Commercial 4.0 International (CC BY-NC 4.0) (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits remixing, adapting and building upon the work for non-commercial use, provided the original work is properly cited.