



Available Online at: www.ijpar.com

[Research article]

SYNTHESIS, ANTI-HIV ACTIVITY AND CYTOTOXICITY OF 2-PHENYL, 3-SULPHONAMIDO QUINAZOLIN-4 (3H) -ONES

¹P.Selvam, C. ²Pannecouque, ²E. De Clercq

¹Raghavendra Institute of Pharmaceutical Education and Research, Anantapur, A.P., India.

²Rega Institute for Medical Research, Katholieke Universiteit-Leuven. Minderbroederstraat 10, Leuven B-3000, Belgium.

ABSTRACT

A series of novel 2,3-disubstitutedquinazolin-4(3H)-ones have been synthesized by condensation of 2-substituted benzo [1,3] oxazine-4-ones and primary amines. Their chemical structures were assigned by means of spectral analysis (FT-IR and ¹H-NMR). Synthesized compounds were screened for *in vitro* antiviral activity against HIV-1 and -2 replication in MT-4 cells. Cytotoxicity was also investigated in uninfected MT-4 cells (C-type Adult T Leukemia cells). All the synthesized compounds exhibited cytotoxicity in MT-4 cells (CC₅₀: 1.93-87 µg/ml) and compound SPB-III displayed marked cytostatic properties in MT-4 cells (CC₅₀: 1.93± 0.29 µg/ml).

Key words: Quinazoline, HIV, MT-4 Cells, MTT assay

INTRODUCTION

Quinazolin-4-(3H)-one is a versatile lead molecule for the design of potential bioactive agents and its derivatives were reported to possess broad spectrum activities. 2-Phenyl -3 - Substituted Quinazolin-4-(3H)-ones were reported to have anti-HIV, some of their derivatives have also shown significant anti-HIV activity^{1,2,3}, anti-cancer activity were studied for 2,3-di substituted quinazolinones derivatives and they showed promising anticancer potential^{4,5,6}. Quinazolinones were screened for their board spectrum anti-viral activity and they were found to be potential derivatives for further studies^{7,8,9}.

Anthranilic acid reaction with benzoyl chloride yields 2-phenyl-1,3-benzoxazin-4-one by N-

acylation via dehydrative cyclization⁸. A series of some novel 2,3-disubstituted quinazolin-4(3H)-one derivatives have been synthesized by condensation of primary aromatic amino group of sulphonamides with 2-substituted-1,3 - benzoxazine-4 - one¹¹ (Scheme-1). These compounds were previously investigated for antiviral activity against bio defence viruses and some of these derivatives inhibits the influenza viruses¹¹. Present work is to investigate the anti-HIV activity and cytotoxicity of Novel 2-phenyl-3-sulphonamido quinazolin-4(3H)-one derivatives (Table 1)

EXPERIMENTAL

Melting points were determined using open ended capillary tube method and are uncorrected. FT-IR

* Corresponding author:

E-mail address: periyasamy_selvam@yahoo.co.in

recorded on Perkin Elmer-1605 series FT-IR in KBr disc. ^1H NMR Spectra was recorded at 400 MHz on Bruker FT-NMR spectrophotometer using TMS as internal standard. Mass spectra was recorded on a Varian Atlas CH-7 Mass Spectrophotometer at 70 eV.

Synthesis of 2-phenyl-3-substituted quinazolin-4-(3H)-one derivatives

An equimolar (0.01 mol) mixture of 2-phenyl-1,3-benzoxazine-4-one and aromatic primary amine sulphonamide derivatives (Sulphanilic acid, sulphanilamide, sulphadiazine, sulphadimidine, sulphamoxole and sulphamethoxazole) was refluxed for 6 hours with 10 ml of acetic acid¹¹. The mixture was cooled to room temperature and poured over crushed ice, filter and then washed with water. The solid thus obtained was recrystallised from ethanol.

Anti-HIV activity

The compounds were tested for anti-HIV activity against the replication of HIV-1(III_B) and HIV-2(ROD) in MT-4 cells⁸. The cells were grown and maintained in RPMI 1640 Medium supplemented with 10% heat-inactivated Fetal Calf Serum (FCS), 2 mM- glutamine, 0.1% Sodium bicarbonate and 20 µg/ml gentamicin (culture medium). HIV-1 (HTLV-III_B/LAI) strain and HIV-2 (LAV-2_{ROD}) strain was used in the experiment. The virus strains were propagated in MT-4 cells. Titer of virus stock was determined in MT-4 cells, and the virus stock was stored at - 70°C until used.

Inhibitory effects of the compounds on HIV-1 and HIV-2 replication were monitored by inhibition of virus-induced cytopathic effect in MT-4 cells and


were estimated by MTT assay. Briefly, 50 µl of HIV-1 and HIV-2 (100-300 CCID₅₀) was added to a flat-bottomed MT-4 cells (6×10^5 cells/ml). After 5 days of incubation, at 37°C the number of viable cells were determined by the 3 - (4, 5-dimethylthiazol-2-yl)-2, 5-diphenyl tetrazolium bromide (MTT) method. Cytotoxicity of the compounds for mock- infected MT-4 cells was assessed by the MTT method. Anti-HIV activity and cytotoxicity of standard AZT were also performed by a similar method in MT-4 cells. The anti-HIV activity and cytotoxicity data are presented in Table 2.


RESULTS

2-phenyl-3-sulphonamido quinazolin-4(3H)-one derivatives were synthesized by condensation of the compounds containing primary aromatic amino group of sulphonamide derivatives with 2-phenyl-1,3-benzoxazin-4-one¹¹. These compounds were previously investigated for antiviral activity against bio-defense viruses and some of these derivatives inhibited the influenza viruses¹¹.

The inhibitory effect of 2-phenyl-3-sulphonamido quinazolin-4 (3H)-ones on the HIV-induced cytopathic effect (CPE) in human lymphocyte MT-4 cell culture was determined by the MT-4/MTT assay. Cytotoxicity was also investigated in uninfected MT-4 cells (C-type Adult T Leukemia cells). All the synthesized compounds exhibited cytotoxicity in MT-4 cells (CC₅₀: 1.93-87 µg/ml) and Compound SPB-III displayed marked cytostatic properties in MT-4 cells (CC₅₀: 1.93 ± 0.29 µg/ml).

Synthesis of 2-Phenyl-3-substituted Quinazolin-4(3H)-one derivatives





COMPOUND CODE	R ₁	R ₂	R-NH ₂
PB-SA	H	H	
PB-SM	H	H	
MBR-SA	Br	H	
MBR-SD	Br	H	
MBR-SDM	Br	H	
MBR-SM	Br	H	
DBR-SD	Br	Br	
DBR-SDM	Br	Br	
PSAVA	H	H	
SPB-III	H	H	
SPF-IIIBr	Br	H	

Table 1: Anti-HIV activity and cytotoxicity of Quinazolin-3(4H)-one derivatives

Compounds	Strain	IC ₅₀ ^a (µg/ml)	CC ₅₀ ^b (µg/ml)	Maximum Protection
PB-SA	IIIB	>67.58	67.58±10.88	16
	ROD	>67.58	67.58±10.88	12
PB-SM	IIIB	>60.88	60.88±10.46	8
	ROD	>60.88	60.88±10.46	1
MBR-SA	IIIB	>87.15	87.15±24.70	3
	ROD	>87.15	87.15±24.70	5
MBR-SD	IIIB	>70.73	70.73±9.17	14
	ROD	>70.73	70.73±9.17	12
MBR-SDM	IIIB	>58.45	58.45±12.18	5
	ROD	>58.45	58.45±12.18	4
MBR-SM	IIIB	>58.08	58.08±11.76	3
	ROD	>58.08	58.08±11.76	5
DBR-SD	IIIB	>72.20	≥72.20	6
	ROD	>72.20	≥72.20	2
DBR-SM	IIIB	>71.70	70.73±19.40	5
	ROD	>71.70	70.73±19.40	8
PSA-VA	IIIB	>89.08	89.08±14.20	4
	ROD	>89.08	89.08±14.20	8
SPB-III	IIIB	>1.93	1.93±0.29	2
	ROD	>1.93	1.93±0.29	1
SPF-IIIBr	IIIB	>96.20	≥96.20	1
	ROD	>96.20	≥96.20	1
SP-II	IIIB	>36.63	36.63±9.14	1
	ROD	>36.63	36.63±9.14	2
S-II	IIIB	>87.20	87.20±22.75	4
	ROD	>87.20	87.20±22.75	5
AZT (STD)	IIIB	0.0015±0.0002	>25	96
Zidovudine	ROD	0.0016±0.0003	>25	76

^aEffective concentration of compound, achieving 50% protection of MT-4 cells against the cytopathic effect of HIV.

^b50% Cytotoxic concentration of compound, required to reduce the viability of mock infected MT-4 cells by 50%.

HIV 1-(HTLV IIIB): HIV-2 (ROD)

REFERENCES

- [1] Shah BR, Bhatt JJ, Patel HH, Undavia NK., Trivedi PB, Desai NC (1995). Synthesis of 2,3-disubstituted-3,1-quinazolin-4(4H)-ones as potential anticancer and anti-HIV agents. *Indian J. Chem.* 34:201-208
- [2] Desai NC, Undavia NK, Trivedi PB & Dipika Dave Vyas GD (1998) Synthesized and screened anti-HIV activity of some non-nucleoside 2,3-disubstitutedquinazoline derivatives *Indian J. Exp. Biol.* 36:1280-1283.
- [3] Alagarsamy V, Pathak US, Pandaya SN, Sriram D & De Clercq E (2000) Anti-HIV and anti bacterial activities of some disubstituted quinazolones and their bio-isoster disubstituted thienopyrimidones. *Indian J. Pharm. Sci.* 66: 433-437.
- [4] Raffa D, Daidone G, Maggio B, Schillaci D, Plescia F (1999). Synthesis and antiproliferative activity of novel 3-(indazol-3-yl)-quinazolin-4(3H) one and 3-(indazol-3-yl)-benzotriazin-4(3H)one derivatives. *Pharmazie* 332: 317-320.
- [5] Murugan V, Padmavathy NP, Ramasarma GVS, Sharma SV, Suresh B (2003). Synthesis of some quinazolinone derivatives as possible anticancer agent. *Indian J. Heterocyclic Chem.* 13: 143-146.
- [6] Girija K, Selvam P & Nagarajan R (2005) Synthesis anticancer activity of 3-[5-Amino-6-(2,3-dichlorophenyl)-[1,2,4]triazin-3-yl]-6,8-dibromo-2-substituted-3H-quinazolin-4-one, *Asian J. Chem.* 17: 1111-1115.
- [7] Manoj K, Srivastava S, Bharati M, Nizamuddin N (2001) Pharmacological studies of some 2-methyl-3-(arythio-carbamido)quinazol-4(3H)-ones and antibacterial activity against *Bacillus cereus*, *S. aureus*, *S. lutea* and antiviral activity against *Gomphrena mosaic*. *Indian J. Chem.* 40: 342-344.
- [8] Selvam P, Dinakaran M, De Clercq E, Sridhar SK (2003). Synthesis, antiviral and cytotoxic activity of 6-bromo-2,3-disubstituted-4(3H)-quinazolinones. *Biological Pharmaceutical Bulletin* 26: 1278-1282.
- [9] Pandey VK, (1996) Synthesized 7-(2'phenyl-3'-ethyl-4'-oxoquinazolinyl)-3,4-diphenyl isoquinolines and screened for antiviral activity against vaccinia virus. *Indian Drugs* 26: 168-171.
- [10] Selvam P, Chennama B, De Clercq E (2004). Synthesis and antiviral activity of some novel 2-substituted 3-(6-ethyl, 4-amino, 5-(4-chlorophenyl)-pyrimidin-2-yl) quinazolin-4(3H)ones. *International J. Chem. Sci.* 2: 627-631
- [11] Selvam P, Vijayalakshimi P, Smee DF, Gowen BB, Julander JG, Day CW, Barnard D L. Novel 3-sulphonamido-quinazolin-4(3H)-one derivatives: microwave-assisted synthesis and evaluation of antiviral activities against respiratory and biodefense viruses. *Antivir Chem Chemother.* 2007; 18(5):301-5.
